

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 672 158 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
17.01.2001 Bulletin 2001/03

(51) Int Cl.7: **C12N 15/87**, A61K 48/00,
C12N 15/34, C12N 15/86,
C12N 15/62, C12N 15/13

(21) Application number: **93924692.2**

(86) International application number:
PCT/GB93/02267

(22) Date of filing: **04.11.1993**

(87) International publication number:
WO 94/10323 (11.05.1994 Gazette 1994/11)

(54) **Adenovirus with genetic fusion of penton fibre to peptide altering binding specificity**

Adenovirus mit genetischer Fusion des Penton Fibers zu einem Peptid das zu einer veränderten
Bindungseigenschaft und Spezifität für Zielzellen führt

Adénovirus comprenant des fibres penton fusionnées à des peptides altérant la spécificité de liaison
pour des cellules cibles.

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**

(56) References cited:
**EP-A- 0 508 809 WO-A-92/06180
WO-A-92/14829 WO-A-93/09221**

(30) Priority: **04.11.1992 GB 9223084**

(43) Date of publication of application:
20.09.1995 Bulletin 1995/38

(60) Divisional application: **00304480.7 / 1 038 967**

(73) Proprietor: **TRANSGENE S.A.**
67082 Strasbourg Cédex (FR)

(72) Inventors:
• **SPOONER, Robert Anthony;**
ICRF Oncology Group
Du Cane Road, London WC12 0HS (GB)
• **EPENETOS, Agamemnon Antoniou;**
ICRF Oncology Group
Du Cane Road, London WC12 0HS (GB)

(74) Representative: **Thomas, Philip John Duval**
Eric Potter Clarkson,
Park View House,
58 The Ropewalk
Nottingham NG1 5DD (GB)

- **NUCLEIC ACIDS RESEARCH.** vol. 21, no. 5, 11 March 1993, ARLINGTON, VIRGINIA US pages 1081 - 1085 **RUSSELL, S.J. ET AL.** 'Retroviral vectors displaying functional antibody fragments' cited in the application
- **HUMAN GENE THERAPY** vol. 3, no. 2, April 1992 pages 147 - 154 **CURIEL, D.T. ET AL.** 'High-efficiency gene transfer mediated by adenovirus coupled to DNA-polylysine complexes' cited in the application
- **TRENDS IN BIOTECHNOLOGY.** vol. 9, September 1991, CAMBRIDGE GB pages 303 - 309 **KINGSMAN, A.J. ET AL.** 'Retroelement particles as purification, presentation and targeting vehicles' cited in the application
- **JOURNAL OF BIOLOGICAL CHEMISTRY.** (MICROFILMS) vol. 268, no. 10, 5 April 1993, BALTIMORE, MD US pages 6866 - 6869 **SHARON, I.M. ET AL.** 'Binding-incompetent adenovirus facilitates molecular conjugate-mediated gene transfer by the receptor-mediated endocytosis pathway' cited in the application

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 672 158 B1

Description

[0001] The present invention relates to delivery vehicles for genes to target cells, especially in the fields of gene therapy and cancer treatment.

[0002] The delivery of genes to target cells, especially those within the mammalian body, has many uses, for example in the fields of gene therapy, cancer treatment and in areas of genetic manipulation still to be discovered. The gene to be delivered may encode a molecule, such as a protein or RNA, which is cytotoxic to the target cell, or it may encode a functional copy of a gene that is defective in the target cell. In this latter case the product of the aforementioned functional copy of the gene will replace that of the defective copy, and the target cell will be able to perform its proper function.

[0003] The use of viruses, or virus-like particles, to deliver genes for gene therapy and cancer treatment has been disclosed.

[0004] However, in most cases the targeting of the virus or virus-like particles containing the desired gene to the cell has relied on the natural host-virus specificity or on local application of the virus to the cells to be targeted, for example direct application of viruses to lung cells by inhalation.

[0005] The human adenovirus 5 (Ad5) genome consists of a double-stranded linear DNA molecule of 36 kilo-base-pair. The virus replication cycle has two phases: an early phase, during which four transcriptional units E1, E2, E3, and E4 are expressed, and a late phase occurring after the onset of viral DNA synthesis when late transcripts are expressed from the major late promoter (MLP). These late messages encode most of the viral structural proteins. E1, E2, and E4 gene products of human adenoviruses (Ads) are involved in transcriptional activation, cell transformation, and viral DNA replication as well as other viral functions, and are essential for viral growth. In contrast, E3 gene products are not required for viral replication in cultured cells or for acute lung infection of cotton rats, but appear to be involved in evading immune surveillance *in vivo*.

[0006] By "virus-like particle" we mean a nucleoprotein particle containing a core of nucleic acid surrounded by protein which (i) is not infective and (ii) can only be propagated in a suitable cell system following transformation by its nucleic acid. Thus a virus-like particle of mammalian origin may be propagated in *Saccharomyces cerevisiae* or in insect cells *via* a baculovirus expression system.

[0007] The modification of coat proteins of filamentous bacteriophages (bacterial viruses), such as M13 and fd, so as to generate novel binding properties, has been disclosed in Cwirla *et al* (1990) *Proc. Natl. Acad. Sci. USA* **87**, 6378-6382 and Scott & Smith (1990) *Science* **249**, 386-390.

[0008] It has previously been suggested that retroelement particles, including retroviral vectors, may be modified to target specific cells, for example see Kingsman *et al* (1991) *Tibtech* **9**, 303-309.

[0009] Russell *et al* (1993) *Nucl. Acids Res.* **21**, 1081-1085, published after the priority date for this application but before the filing date discloses retroviral vectors displaying functional antibody fragments and suggests that, in principle, the display of antibody fragments on the surface of recombinant retroviral particles could be used to target virus to cells for gene delivery. However, it is not known whether a retrovirus can be assembled in which all the subunits of the viral envelope protein are fused to antibody, and if so whether the virus would infect cells.

[0010] NIP-derivatised human cells were tested as a method for targeted gene delivery, but became permissive for both modified (displaying an anti-NIP antibody) and unmodified ecotropic viral particles. NIP is 4-hydroxy-3-iodo-5-nitrophenylacetic acid.

[0011] Michael *et al* (1993) *J. Biol. Chem.* **268**, 6866-6869, published after the priority date of this application but before the filing date, describes molecular conjugates between adenovirus and a vector system comprising two linked domains, a DNA binding domain and a ligand domain. In this configuration, however, it is stated that the viral moiety functions in the capacity of both an alternate ligand domain of the conjugate and, since an additional ligand has been introduced into the conjugate design, the potential for cell-specific targeting is undermined.

[0012] Curiel *et al* (1992) *Human Gene Therapy* **3**, 147-154 describes adenoviruses wherein a foreign epitope was introduced into the hexon protein and polylysine-antibody complexed DNA was attached to adenovirus by virtue of the antibody binding the foreign epitope on the hexon. Foreign DNA is transferred bound to the exterior of the virion.

[0013] WO 92/06180 discusses in general terms adding a binding moiety to the surface of a virus and gives examples of modified Moloney murine leukaemia virus and hepatitis B virus.

[0014] WO 92/14829 discusses that specific membrane-associated protein from an unrelated virus can be incorporated into retroviral particles to provide an altered host range.

[0015] EP 0 508 809 describes a lipid-containing virus-like particle based on a retroelement such as a retrovirus.

[0016] WO 93/09221 discusses in general terms modifications to a cell-binding receptor of a virus. Epstein-Barr Virus, Influenza A virus, paramyxovirus and retroviruses are specifically mentioned.

[0017] The above-mentioned viruses and virus-like particles may be able to target cells using the binding moiety displayed on their surface but they can also still target their natural host cells.

[0018] We have now devised new viruses and virus-like particles at least some of which can bind the target cell with

high specificity and may deliver genetic material to the target cell; at least some of the viruses and virus-like particles may bind and deliver genetic material to the target cell without substantially binding to the natural host cell of the virus.

[0019] One aspect of the present invention provides an adenovirus, or adenovirus-like particle, or a replication defective derivative thereof comprising a modified binding specificity conferred by a binding moiety allowing the adenovirus or adenovirus-like particle to bind to a target cell, wherein the binding moiety is a peptide and is fused to the penton fibre of the adenovirus or adenovirus-like particle.

[0020] The virus or virus-like particle may be substantially incapable of binding its host cell.

[0021] By "substantially incapable of binding its host cell" we mean that the modified virus has no more than 1 % of the binding affinity of the unmodified virus for the host cell.

[0022] In general, the binding specificity of a natural virus or virus-like particle is conferred by the specific interaction between a receptor-like molecule expressed on the surface of the virus or virus-like particle and a cognate receptor-like molecule expressed on the surface of its host cell. The invention provides a beneficial modification of the binding specificity, so that the virus or virus-like particle can bind to a different specific target cell.

[0023] The introduction of the modified binding moiety may be such as to achieve the said removal of the native binding specificity.

[0024] By "binding moiety" we mean a molecule that is exposed on the surface of the virus or virus-like particle which is able to bind to a molecule on the target cell. The "binding moiety" may be a molecule on the virus or virus-like particle modified in such a way that its binding specificity is changed, or it may be a molecule added to, and exposed on the surface of, the virus or virus-like particle to provide a new binding specificity.

[0025] It is preferred if the binding moiety is external to the receptor for its host cell of the naive, unmodified virus.

[0026] It is further preferred if the binding moiety is joined or fused to the virus or virus-like particles directly or indirectly by a spacer group.

[0027] By "host cell" we mean the cell that an unmodified, naive virus can bind to using its receptor-like molecule and the cognate receptor-like molecule on the cell. By "target cell" we mean the cell that the modified virus can bind to using its binding moiety. In some circumstances in the context of the second aspect of the invention, such as when the binding moiety recognises an entity on the host cell which is not the cognate receptor-like molecule, then the host cell may be the target cell.

[0028] It is expected that the invention will find uses in the areas of gene therapy and cancer treatment.

[0029] It is also preferred that the virus or virus-like particle is "replication-defective". By "replication defective" we mean a virus whose genetic material has been manipulated so that it cannot divide or proliferate in the cell it infects.

[0030] The binding moiety of the virus or virus-like particle of the invention provides the target cell binding specificity. Any cell-binding protein or peptide may be useful for targeting the virus or virus-like particle to the cell. For example, short linear stretches of amino acids, such as those constituting a peptide hormone, are useful, as are domains of polypeptides that can fold independently into a structure that can bind to the target cell.

[0031] In one preferred embodiment the binding moiety has the property of any one of a monoclonal antibody, ScFv (single chain Fv fragment), a dAb (single domain antibody) or a minimal recognition unit of an antibody.

[0032] The binding site on the target cell may be a target cell-specific antigen. Such antigens are listed in Table 1. Other binding moieties, targets on cells, and diseases which could usefully be treated using reagents delivered by the modified viruses or virus-like particles are given in Table 2.

Table 1

1. Tumour Associated Antigens		
Antigen	Antibody	Existing Uses
Carcino-embryonic Antigen	{C46 (Amersham) {85A12 (Unipath)	Imaging & Therapy of colon/rectum tumours.
Placental Alkaline Phosphatase	H17E2 (ICRF, Travers & Bodmer)	Imaging & Therapy of testicular and ovarian cancers.
Pan Carcinoma	NR-LU-10 (NeoRx Corporation)	Imaging & Therapy of various carcinomas incl. small cell lung cancer.
Polymorphic Epithelial Mucin (Human milk fat globule)	HMFG1 (Taylor-Papadimitriou, ICRF)	Imaging & Therapy of ovarian cancer, pleural effusions.

Table 1 (continued)

1. Tumour Associated Antigens		
Antigen	Antibody	Existing Uses
β -human Chorionic Gonadotropin	W14	Targeting of enzyme (CPG2) to human xenograft choriocarcinoma in nude mice. (Searle <i>et al</i> (1981) <i>Br. J. Cancer</i> 44 , 137-144)
A Carbohydrate on Human Carcinomas	L6 (IgG2a) ¹	Targeting of alkaline phosphatase. (Senter <i>et al</i> (1988) <i>P.N.A.S.</i> 85 , 4842-4846)
CD20 Antigen on B Lymphoma (normal and neoplastic)	1F5 (IgG2a) ²	Targeting of alkaline phosphatase. (Senter <i>et al</i> (1988) <i>P.N.A.S.</i> 85 , 4842-4846)
Other antigens include alphafoetoprotein, Ca-125 and prostate specific antigen.		
2. Immune Cell Antigens		
Pan T Lymphocyte Surface Antigen (CD3)	OKT-3 (Ortho)	As anti-rejection therapy for kidney transplants.
B-lymphocyte Surface Antigen (CD22)	RFB4 (Janossy, Royal Free Hospital)	Immunotoxin therapy of B cell lymphoma.
Pan T lymphocyte Surface Antigen (CD5)	H65 (Bodmer, Knowles ICRF, Licensed to Xoma Corp., USA)	Immunotoxin treatment of Acute Graft versus Host disease, Rheumatoid Arthritis.
3. Infectious Agent-Related Antigens		
Mumps virus-related	Anti-mumps polyclonal antibody	Antibody conjugated to Diphtheria toxin for treatment of mumps.
Hepatitis B Surface Antigen	Anti HBs Ag	Immunotoxin against Hepatoma.

¹Hellström *et al* (1986) *Cancer Res.* **46**, 3917-3923²Clarke *et al* (1985) *P.N.A.S.* **82**, 1766-1770

Table 2:

Binding moieties for tumour-specific targets and tumour associated antigens		
Target	Binding moiety	Disease
Truncated EGFR Idiotypes EGFR (<i>c-erbB1</i>)	anti-EGFR mAb anti-id mAbs EGF, TGF α anti-EGFR mAb	Gliomas B-cell lymphomas Breast cancer
<i>c-erbB2</i>	mAbs	Breast cancer
IL-2 receptor	IL-2	Lymphomas and leukaemias
	anti-Tac mAb	
IL-4 receptor	IL-4	Lymphomas and leukaemias
IL-6 receptor	IL-6	Lymphomas and leukaemias
MSH (melanocyte-stimulating hormone) receptor	α -MSH	Melanomas
Transferrin receptor (TR)	Transferrin anti-TR mAb	Gliomas
gp95/gp97	mAbs	Melanomas

Table 2: (continued)

Binding moieties for tumour-specific targets and tumour associated antigens		
Target	Binding moiety	Disease
p-glycoprotein cells	mAbs	drug-resistant
cluster-1 antigen (N-CAM)	mAbs	Small cell lung carcinomas
cluster-w4	mAbs	Small cell lung carcinomas
cluster-5A	mAbs	Small cell lung carcinomas
cluster-6 (LeY)	mAbs	Small cell lung carcinomas
PLAP (placental alkaline phosphatase)	mAbs	Some seminomas Some ovarian; some non-small cell lung cancer
CA-125	mAbs	Lung, ovarian
ESA (epithelial specific antigen)	mAbs	carcinoma
CD 19, 22, 37	mAbs	B-cell lymphoma
250 kDa proteoglycan	mAbs	Melanoma
p55	mAbs	Breast cancer
TCR-IgH fusion	mAbs	Childhood T-cell leukaemia
Blood gp A antigen (in B or O individuals)	mAbs	Gastric and colon tumours

[0033] The binding moiety may be a monoclonal antibody. Monoclonal antibodies which will bind to many of these antigens are already known but in any case, with today's techniques in relation to monoclonal antibody technology, antibodies can be prepared to most antigens. The binding moiety may be a part of an antibody (for example a Fab fragment) or a synthetic antibody fragment (for example, ScFv). Suitable monoclonal antibodies to selected antigens may be prepared by known techniques, for example those disclosed in *"Monoclonal Antibodies: A manual of techniques"*, H Zola (CRC Press, 1988) and in *"Monoclonal Hybridoma Antibodies: Techniques and Applications"*, J G R Hurrell (CRC Press, 1982).

[0034] Suitably prepared non-human antibodies can be "humanized" in known ways, for example by inserting the CDR regions of mouse antibodies into the framework of human antibodies.

[0035] The variable heavy (V_H) and variable light (V_L) domains of the antibody are involved in antigen recognition, a fact first recognised by early protease digestion experiments. Further confirmation was found by "humanization" of rodent antibodies. Variable domains of rodent origin may be fused to constant domains of human origin such that the resultant antibody retains the antigenic specificity of the rodent parental antibody (Morrison *et al* (1984) *Proc. Natl. Acad. Sci. USA* **81**, 6851-6855).

[0036] That antigenic specificity is conferred by variable domains and is independent of the constant domains is known from experiments involving the bacterial expression of antibody fragments, all containing one or more variable domains. These molecules include Fab-like molecules (Better *et al* (1988) *Science* **240**, 1041); Fv molecules (Skerra *et al* (1988) *Science* **240**, 1038); ScFv molecules where the V_H and V_L partner domains are linked via a flexible oligopeptide (Bird *et al* (1988) *Science* **242**, 423; Huston *et al* (1988) *Proc. Natl. Acad. Sci. USA* **85**, 5879) and dAbs comprising isolated V domains (Ward *et al* (1989) *Nature* **341**, 544). A general review of the techniques involved in the synthesis of antibody fragments which retain their specific binding sites is to be found in Winter & Milstein (1991) *Nature* **349**, 293-299.

[0037] By "ScFv molecules" we mean molecules wherein the V_H and V_L partner domains are linked via a flexible oligopeptide.

[0038] It may be advantageous to use antibody fragments, rather than whole antibodies. Effector functions of whole antibodies, such as complement binding, are removed. ScFv and dAb antibody fragments can be expressed as fusions with other polypeptides.

[0039] Minimal recognition units may be derived from the sequence of one or more of the complementary-determining regions (CDR) of the Fv fragment. Whole antibodies, and $F(ab')_2$ fragments are "bivalent". By "bivalent" we mean that the said antibodies and $F(ab')_2$ fragments have two antigen combining sites. In contrast, Fab, Fv, ScFv, dAb fragments and minimal recognition units are monovalent, having only one antigen combining sites.

[0040] In a further embodiment the binding moiety is at least part of a ligand of a target cell-specific cell-surface receptor.

[0041] It is preferred that the target cell-specific cell-surface receptor is the receptor for human gonadotrophin-releasing hormone (GnRH). In this preferred embodiment the binding moiety is GnRH, and its binding specificity is for human cancer cells that express the GnRH receptors on their surface. Examples of such human cancer cells are

prostate, breast and endometrial cancer cells.

[0042] It is also preferred that the target cell-specific cell-surface receptor is the receptor for melanocyte-stimulating hormone (MSH) which is expressed in high number in melanoma cells. In this preferred embodiment the binding moiety is MSH, and its binding specificity is for melanoma cells.

[0043] It is also preferred that the target cell-specific cell-surface receptor is the receptor for somatostatin.

[0044] Of course, the receptors for GnRH, MSH and somatostatin may themselves be target cell-specific antigens and may be recognised by binding moieties which have the property of any one of a monoclonal antibody, a ScFv, a dAb or a minimal recognition unit. Thus, although the binding site on the target cell may be a cell-surface receptor it may also act as a target cell-specific cell-surface antigen for recognition by the binding moiety.

[0045] It will be appreciated by those skilled in the art that binding moieties which are polypeptides may be conveniently made using recombinant DNA techniques. The binding moiety may be fused to a protein on the surface of the virus or virus-like protein as disclosed below or they may be synthesised independently of the virus or virus-like particle, by expression from a suitable vector in a suitable host and then joined to the virus or virus-like particle as disclosed below.

[0046] Nucleic acid sequences encoding many of the targeting moieties are known, for example those for peptide hormones, growth factors, cytokines and the like and may readily be found by reference to publicly accessible nucleotide sequence databases such as EMBL and GenBank. Once the nucleotide sequence is known it is obvious to the person skilled in the art how to make DNA encoding the chosen binding moiety using, for example, chemical DNA synthetic techniques or by using the polymerase chain reaction to amplify the required DNA from genomic DNA or from tissue-specific cDNA.

[0047] Many cDNAs encoding peptide hormones, growth factors, cytokines and the like, all of which may be useful as binding moieties, are generally available from, for example British Biotechnology Ltd, Oxford, UK.

[0048] It is preferred that when the virus or virus-like particle of the invention binds to its target cell it delivers its nucleic acid to the said target cell, that is the target cell is infected by the virus or virus-like particle. Target cells, especially cancer cells, that are infected in this manner by the virus or virus-like particle may express viral molecules on their surface and may be recognised by the immune system and destroyed. Of course, other cytotoxic functions of the virus may also kill the cell.

[0049] In one embodiment, the adenovirus E1B gene is substantially deleted or modified so that its gene product no longer interacts with the E1A protein. E1A protein stimulates apoptosis but normally its action is inhibited by E1B. Conveniently, the E1B gene is inactivated by insertion; preferably a cytotoxic gene, as defined below, is inserted at or near the E1B gene.

[0050] E1, E3 and a site upstream of E4 may be used as sites for insertion of foreign DNA sequences in the generation of recombinant adenoviruses for example see Berkner and Sharp (1984) *Nucl. Acids Res.* **12**, 1925-1941; Chanda *et al* (1990) *Virology* **175**, 535-547; Haj-Ahmad and Graham (1986) *J. Virol.* **57**, 267-274; Saito *et al* (1985) *J. Virol.* **54**, 711-719; all incorporated herein by reference. Since the upper size limit for DNA molecules that can be packaged into adenovirus particles is approximately 105% of the wild-type genome only about 2 kb of extra DNA can be inserted without compensating deletions of viral DNA. Although E1 is essential for virus replication in cell culture, foreign DNA can be substituted for E1 sequences when the virus is grown in 293 cells which are transformed by Ad5 DNA and constitutively express E1 (Graham *et al* (1977) *J. Gen. Virol.* **36**, 59-72, incorporated herein by reference). Several vectors having 1.9 kb deleted from E3 of Ad5 have been constructed without interfering with virus replication in cell culture (reviewed by Graham and Prevec (1992) in "Vaccines: New Approaches to Immunological Problems" R.W. Ellis (Ed.), Butterworth-Heinemann, Boston, MA, pages 364-390, incorporated herein by reference). Such vectors allow for insertion of up to 4 kb of foreign DNA. Recombinant adenoviruses containing inserts in E3 replicate in all Ad-permissive cell lines and a number of adenovirus vectors containing E3 inserts have been shown to express foreign genes efficiently both *in vitro* and *in vivo* (Berkner (1988) *Biotechniques* **6**, 616-629; Chanda *et al* (1990) *Virology* **175**, 535-547; Dewar *et al* (1989) *J. Virol.* **63**, 129-136; Graham (1990) *Trends Biotechnol.* **8**, 85-87; Graham and Prevec (1992) in "Vaccines: New Approaches to Immunological Problems" R.W. Ellis (Ed.), Butterworth-Heinemann, Boston, MA, pages 364-390; Johnson *et al* (1988) *Virology* **164**, 1-14; Lubeck *et al* (1989) *Proc. Natl. Acad. Sci. USA* **86**, 6763-6767; McDermott *et al* (1989) *Virology* **169**, 244-247; Morin *et al* (1987) *Proc. Natl. Acad. Sci. USA* **84**, 4626-4630; Prevec *et al* (1989) *J. Gen. Virol.* **70**, 429-434; Prevec *et al* (1990) *J. Inf. Dis.* **161**, 27-30; Schneider *et al* (1989) *J. Gen. Virol.* **70**, 417-427; Vernon *et al* (1991) *J. Gen. Virol.* **72**, 1243-1251; Yuasa *et al* (1991) *J. Gen. Virol.* **72**, 1927-1934) all incorporated herein by reference.

[0051] Substantially replication-defective adenoviruses may be made by creating a deficiency of the E1A protein. Suitably this is achieved by deleting the E1A gene or by making mutations within the E1A gene that prevent expression of the E1A protein. Examples of suitable mutations are deletions within the E1A coding region; nonsense mutations; and frameshift mutations.

[0052] In further preference, the virus or virus-like particle is modified further to contain a gene suitable for gene therapy.

[0053] In one embodiment, the gene encodes a molecule having a directly or indirectly cytotoxic function. By "directly or indirectly" cytotoxic, we mean that the molecule encoded by the gene may itself be toxic (for example ricin; tumour necrosis factor; interleukin-2; interferon-gamma; ribonuclease; deoxyribonuclease; *Pseudomonas* exotoxin A) or it may be metabolised to form a toxic product, or it may act on something else to form a toxic product. The sequence of ricin cDNA is disclosed in Lamb *et al* (1985) *Eur. J. Biochem.* **148**, 265-270 incorporated herein by reference.

[0054] For example, it would be desirable to target a DNA sequence encoding an enzyme using the virus or virus-like particle of the invention, the enzyme being one that converts a relatively non-toxic prodrug to a toxic drug. The enzyme cytosine deaminase converts 5-fluorocytosine (5FC) to 5-fluorouracil (5FU) (Mullen *et al* (1922) *PNAS* **89**, 33); the herpes simplex enzyme thymidine kinase sensitises cells to treatment with the antiviral agent ganciclovir (GCV) or aciclovir (Moolten (1986) *Cancer Res.* **46**, 5276; Ezzedine *et al* (1991) *New Biol* **3**, 608). The cytosine deaminase of any organism, for example *E. coli* or *Saccharomyces cerevisiae*, may be used.

[0055] Thus, in a preferred embodiment of the invention, the gene encodes a cytosine deaminase and the patient is concomitantly given 5FC. By "concomitantly", we mean that the 5FC is administered at such a time, in relation to the transformation of the tumour cells, that 5FC is converted into 5FU in the target cells by the cytosine deaminase expressed from the said gene. A dosage of approximately 0.001 to 100.0 mg 5FC/kg body weight/day, preferably 0.1 to 10.0 mg/kg/day is suitable.

[0056] Components, such as 5FC, which are converted from a relatively non-toxic form into a cytotoxic form by the action of an enzyme are termed "pro-drugs".

[0057] Other examples of pro-drug/enzyme combinations include those disclosed by Bagshawe *et al* (WO 88/07378), namely various alkylating agents and the *Pseudomonas* spp. CPG2 enzyme, and those disclosed by Epenetos & Rowlinson-Busza (WO 91/11201), namely cyanogenic pro-drugs (for example amygdalin) and plant-derived β -glucosidases.

[0058] Enzymes that are useful in this embodiment of the invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine deaminase useful for converting non-toxic 5-fluorocytosine into the anti-cancer drug, 5-fluorouracil; proteases, such as serratio protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; D-alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as β -galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs; β -lactamase useful for converting drugs derivatized with β -lactams into free drugs; and penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxycetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as abzymes, can be used to convert the prodrugs of the invention into free active drugs [see, e.g. R J Massey, *Nature*, **328**, pp. 457-458 (1987)].

[0059] Similarly, the prodrugs of this invention include, but are not limited to, the above-listed prodrugs, e.g., phosphate-containing prodrugs, thiophosphate-containing prodrugs, sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, β -lactam-containing prodrugs, optionally substituted phenoxycetamide-containing prodrugs or optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5-fluorouridine prodrugs which can be converted by the enzyme of the conjugate into the more active, cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, etoposide, teniposide, adriamycin, daunomycin, carminomycin, aminopterin, dactinomycin, mitomycins, cis-platinum and cis-platinum analogues, bleomycins, esperamicins [see U.S. Pat. No. 4,675,187], 5-fluorouracil, melphalan and other related nitrogen mustards.

[0060] In a further embodiment the gene delivered to the target cell encodes a ribozyme capable of cleaving targeted RNA or DNA. The targeted RNA or DNA to be cleaved may be RNA or DNA which is essential to the function of the cell and cleavage thereof results in cell death or the RNA or DNA to be cleaved may be RNA or DNA which encodes an undesirable protein, for example an oncogene product, and cleavage of this RNA or DNA may prevent the cell from becoming cancerous.

[0061] Ribozymes which may be encoded in the genomes of the viruses or virus-like particles herein disclosed are described in Cech and Herschlag "Site-specific cleavage of single stranded DNA" US 5,180,818; Altman *et al* "Cleavage of targeted RNA by RNase P" US 5,168,053, Cantin *et al* "Ribozyme cleavage of HIV-1 RNA" US 5,149,796; Cech *et al* "RNA ribozyme restriction endoribonucleases and methods", US 5,116,742; Been *et al* "RNA ribozyme polymerases, dephosphorylases, restriction endonucleases and methods", US 5,093,246; and Been *et al* "RNA ribozyme polymerases, dephosphorylases, restriction endoribonucleases and methods; cleaves single-stranded RNA at specific site by transesterification", US 4,987,071, all incorporated herein by reference.

[0062] In a still further embodiment the gene delivered to the target cell encodes an antisense RNA.

[0063] By "antisense RNA" we mean an RNA molecule which hybridises to, and interferes with the expression from a mRNA molecule encoding a protein or to another RNA molecule within the cell such as pre-mRNA or tRNA or rRNA,

or hybridises to, and interferes with the expression from a gene.

[0064] Conveniently, a gene expressing an antisense RNA may be constructed by inserting a coding sequence encoding a protein adjacent a promoter in the appropriate orientation such that the RNA complementary to mRNA. Suitably, the antisense RNA blocks expression of undesirable polypeptides such as oncogenes, for example *ras*, *bcl*, *src* or tumour suppressor genes such as p53 and *Rb*.

[0065] It will be appreciated that it may be sufficient to reduce expression of the undesirable polypeptide rather than abolish the expression.

[0066] It will be further appreciated that DNA sequences suitable for expressing as antisense RNA may be readily derived from publicly accessible databases such as GenBank and EMBL.

[0067] In another embodiment of the invention, the gene replaces the function of a defective gene in the target cell.

[0068] There are several thousand inherited genetic diseases of mammals, including humans, that are caused by defective genes. Examples of such genetic diseases include cystic fibrosis, where there is known to be a mutation in the CFTR gene; Duchenne muscular dystrophy, where there is known to be a mutation in the dystrophin gene; sickle cell disease, where there is known to be a mutation in the HbA gene. Many types of cancer are caused by defective genes, especially protooncogenes, and tumour-suppressor genes that have undergone mutation.

[0069] Thus, it is preferred that the virus or virus-like particle of the invention, which may be useful in the treatment of cystic fibrosis, contains a functional CFTR gene to replace the function of the defective CFTR gene. Similarly, it is preferred that the virus or virus-like particle of the invention, which may be useful in the treatment of cancer, contains a functional protooncogene, or tumour-suppressor gene to replace the function of the defective protooncogene or tumour-suppressor gene.

[0070] Examples of protooncogenes are *ras*, *src*, *bcl* and so on; examples of tumour-suppressor genes are p53 and *Rb*.

[0071] By "gene" we mean a nucleic acid coding sequence that may contain introns, or fragment thereof, or cDNA, or fragment thereof.

[0072] It will be appreciated that the gene will be introduced into a convenient place within the genome of the virus or virus-like particle and will contain a promoter and/or enhancer element to drive its expression.

[0073] It is preferred if the promoter and/or enhancer is selective for the cells to be targeted. Some examples of tissue or tumour specific promoters are given below but new ones are being discovered all of the time which will be useful in this embodiment of the invention.

[0074] The tyrosinase and TRP-1 genes both encode proteins which play key roles in the synthesis of the pigment melanin, a specific product of melanocytic cells. The 5' ends of the tyrosinase and tyrosinase-related protein (TRP-1) genes confer tissue specificity of expression on genes cloned downstream of these promoter elements.

[0075] The 5' sequences of these genes are described in Bradl, M. *et al* (1991) *Proc. Natl. Acad. Sci. USA* **88**, 164-168 and Jackson, I.J. *et al* (1991) *Nucleic Acids Res.* **19**, 3799-3804.

[0076] Prostate-specific antigen (PSA) is one of the major protein constituents of the human prostate secretion. It has become a useful marker for the detection and monitoring of prostate cancer. The gene encoding PSA and its promoter region which directs the prostate-specific expression of PSA have been described (Lundwall (1989) *Biochem. Biophys. Res. Comm.* **161**, 1151-1159; Riegman *et al* (1989) *Biochem. Biophys. Res. Comm.* **159**, 95-102; Brawer (1991) *Acta Oncol.* **30**, 161-168).

[0077] Carcinoembryonic antigen (CEA) is a widely used tumour marker, especially in the surveillance of colonic cancer patients. Although CEA is also present in some normal tissues, it is apparently expressed at higher levels in tumorous tissues than in corresponding normal tissues. The complete gene encoding CEA has been cloned and its promoter region analysed. A CEA gene promoter construct, containing approximately 400 nucleotides upstream from the translational start, showed nine times higher activity in the adenocarcinoma cell line SW303, compared with the HeLa cell line. This indicates that cis-acting sequences which convey cell type specific expression are contained within this region (Schrewe *et al* (1990) *Mol. Cell. Biol.* **10**, 2738-2748).

[0078] The *c-erbB-2* gene and promoter have been characterised previously and the gene product has been shown to be over-expressed in tumour cell lines (Kraus *et al* (1987) *EMBO J.* **6**, 605-610).

[0079] The mucin gene, MUC1, contains 5' flanking sequences which are able to direct expression selectively in breast and pancreatic cell lines, but not in non-epithelial cell lines as taught in WO 91/09867.

[0080] The binding moiety allowing the virus or virus-like particle to bind to a target cell may be a polypeptide capable of binding specifically to the target cell.

[0081] It is preferred that the binding moiety is a polypeptide.

[0082] It is preferred that the binding moiety and the molecule on the surface of the virus or virus-like particle are both polypeptides that may be produced as a fusion by the techniques of genetic engineering. The use of genetic engineering allows for the precise control over the fusion of such polypeptides.

[0083] Thus a further embodiment of the invention is a nucleotide sequence encoding the fusion of the binding moiety and the protein on the surface of the virus or virus-like particle.

[0084] The nucleotide sequence encoding the fusion of the binding moiety and the protein on the surface of the virus or virus-like particle is preferably made by an alteration of the viral genome.

[0085] The nucleotide sequence may be synthesised *de novo* using solid phase phosphoramidite chemistry, but it is more usual for the nucleotide sequence to be constructed from two parts, the first encoding the binding moiety and the second the protein on the surface of the virus or virus-like particle. The two parts may be derived from their respective genes by restriction endonuclease digestion or by other methods known by those skilled in the art such as the polymerase chain reaction.

[0086] A variety of methods have been developed to operatively link two nucleotide sequences via complementary cohesive termini. For instance, synthetic linkers containing one or more restriction sites provide a method of joining the two DNA segment together. Each DNA segment, generated by endonuclease restriction digestion, is treated with bacteriophage T4 DNA polymerase or *E. coli* DNA polymerase I, enzymes that remove protruding, 3'-single-stranded termini with their 3'-5'-exonucleolytic activities, and fill in recessed 3'-ends with their polymerizing activities.

[0087] The combination of these activities therefore generates blunt-ended DNA segments. The blunt-ended segments are then incubated with a large molar excess of linker molecules in the presence of an enzyme that is able to catalyze the ligation of blunt-ended DNA molecules, such as bacteriophage T4 DNA ligase. Thus, the products of the reaction are DNA segments carrying polymeric linker sequences at their ends. These DNA segments are then cleaved with the appropriate restriction enzyme and lifted to an expression vector that has been cleaved with an enzyme that produces termini compatible with those of the DNA segment.

[0088] Synthetic linkers containing a variety of restriction endonuclease sites are commercially available from a number of sources including International Biotechnologies Inc, New Haven, CN, USA.

[0089] A desirable way to generate the DNA encoding the fusion polypeptide of the invention is to use the polymerase chain reaction as disclosed by Sails *et al* (1988) *Science* **239**, 487-491.

[0090] In this method each of the DNA molecules encoding the two polypeptides to be fused are enzymatically amplified using two specific oligonucleotide primers which themselves become incorporated into the amplified DNA. The said specific primers may contain restriction endonuclease recognition sites which may then be used to join the said two DNA molecules using T4 DNA ligase as disclosed.

[0091] A particular feature of one aspect of the present invention is the modification of the virus or virus-like particle of the invention so that it no longer binds its host cell and so that it binds the target cell by virtue of its binding moiety.

[0092] The penton fibre may be modified by the insertion or deletion or substitution of amino acid residues that disrupt their host-cell binding function. It is preferred that the binding moiety for the target cell is joined to the host-cell receptor in such a manner that the binding moiety is capable of binding the target cell, the host-cell receptor is unable to bind to the host cell and therefore the binding specificity of the virus or virus-like particle is modified. A further preference is that the portion of the penton fibre that is exposed on the surface of the virus or virus-like particle is replaced by the binding moiety, and that the portion of the penton fibre which promotes the uptake of viral DNA by the target cell is retained. Suitably, the binding moiety is joined directly or indirectly to the host-cell receptor by a spacer group.

[0093] Examples of spacer groups are polypeptide sequences of between 4 and 1000 amino acid residues.

[0094] Thus, in one embodiment of the invention the gene encoding the penton fibre in adenovirus is modified in such a way that the DNA encoding the surface-exposed portion is replaced by a DNA fragment encoding a ScFv, the ScFv being derived from an antibody which binds to a target cell surface antigen.

[0095] Potential fusion sites within the penton fibre have been identified.

[0096] The adenovirus fibre is a trimer composed of three protomers. The amino terminal end (40 amino acids or so) of each participates in the formation of a tail that is closely associated with the penton (as opposed to the hexon) subunit of the capsid. High amino acid conservation is maintained between the different characterised serotypes.

[0097] Middle portions of each protomer form the shaft of the protein. This shaft is of variable length, depending upon serotype, and is composed of repeating units of 15 amino acids (for examples, serotypes have been identified with 6, 15 and 21 repeat units). These repeating units are not duplicates: rather than strict conservation of amino acid structure, there is a general conservation of relative hydrophobicity. Some serotypes, for example, 40 and 41, have shafts composed of different length fibre proteins. This suggests a certain flexibility in structural constraints.

[0098] The carboxy-terminal ends (some 200 amino acids) associate to form a knob that is held erect a great distance (in molecular terms) from the capsid.

[0099] Whilst the cellular receptor(s) and mechanisms of docking have not been firmly identified and elucidated, we propose that the most likely candidate structure for cell binding is the knob. Thus, in one embodiment the whole knob of the penton fibre has been replaced with single chain antibody (ScFv) domains. The triplex structure implies that each fibre will thus end in three ScFvs. Additionally, the ScFv regions can be replaced with CDRs, or by non-antibody derived peptides, of known specificity or other molecules that are capable of interacting specifically with the target cell.

[0100] Suitable fusion sites are therefore at the native junction between shaft and knob domains, or (should the DNA sequence prove to be more amenable) at any junction between repetitive units of the shaft. Preferably, the minimum shaft length is not reduced beyond the smallest size naturally identified. There are thus at least 15 potential sites at

which fusion could be contemplated.

[0101] Although it is preferred that the binding moiety forms the end of the fibre thereby replacing the knob, the binding moiety may also be fused within the penton fibre sequence but still display its binding surfaces and bind to the target cell.

[0102] Suitably, the binding moiety may be fused to the knob and extend externally to the knob structure.

[0103] A further aspect of the invention provides a method of producing in cell culture a virus or virus-like particle and then joining the binding moiety, as defined above, to the virus or virus-like particle.

[0104] A further aspect of the invention provides a method of producing in cell culture a virus or virus-like particle which has been genetically modified to express a binding moiety on its surface. The virus or virus-like particle is grown in its host prior to modification, but once the modification that alters the binding specificity is made, the virus or virus-like particle is grown in the target cell. Thus, for example in the case where the binding moiety recognises a breast tumour cell antigen, the virus or virus-like particle is grown in breast tumour cell culture.

[0105] The virus or virus-like particles of the invention are administered in any suitable way, usually parenterally, for example intravenously, intraperitoneally or intravesically, in standard sterile, non-pyrogenic formulations of diluents and carriers, for example isotonic saline (when administered intravenously).

[0106] A further aspect of the invention provides a method of delivery of the virus or virus-like particle which contains a gene encoding a molecule having an indirectly cytotoxic function.

[0107] Suitably, the indirectly cytotoxic function is an enzyme that converts a prodrug to a toxic drug. With such a virus or virus-like particle, once the virus or virus-like particle has bound to the target cells, delivered its nucleic acid to the cells, and expressed the indirectly cytotoxic functions, which typically takes a day or so, the pro-drug is administered. The timing between administration of the virus or virus-like particle and the pro-drug may be optimised in a non-inventive way.

[0108] The dosage of the pro-drug will be chosen by the physician according to the usual criteria. The dosage of the virus or virus-like particle will similarly be chosen according to normal criteria, and in the case of tumour treatment, particularly with reference to the type, stage and location of tumour and the weight of the patient. The duration of treatment will depend in part upon the rapidity and extent of any immune reaction to the virus or virus-like particle.

[0109] Some of the viruses or virus-like particles either in themselves, or together with an appropriate pro-drug, are in principle suitable for the destruction of cells in any tumour or other defined class of cells selectively exhibiting a recognisable (surface) entity. Examples of types of cancer that may be treated using the viruses or virus-like particles are cancer of the breast, prostate, colon, rectum, ovary, testicle and brain. The compounds are principally intended for human use but could be used for treating other mammals including dogs, cats, cattle, horses, pigs and sheep.

[0110] The invention will now be described in detail with reference to the following Figures and Examples in which:

[0111] Figure 1 shows (a) an unmodified (i.e. "naive") virus or virus-like particle able to bind to and infect its host cell but not a non-host cell, such as a target cell; and (b) a virus or virus-like particle with a modified binding specificity does not bind and infect its host cell but binds and infects a target cell; and (c) a virus or virus-like particle as in (b) modified further to contain a gene for gene therapy or cancer treatment.

[0112] Figure 2 shows (a) unmodified (naive) adenovirus; (b) adenovirus modified so that its penton fibres, which recognise the host cell, are replaced in part by antibody fragments which recognise the target cell; and (c) adenovirus as in (b) with further genetic material added to the viral DNA for gene therapy of cancer.

[0113] Figure 3 shows (a) influenza virus and (b) genetically-modified influenza virus wherein at least part of the haemagglutinin binding site is replaced by an antibody with anti-cancer cell binding activity. The virus shown in Figure 3 does not fall within the scope of the claims.

[0114] Figure 4 shows (a) a retrovirus virus; and (b) as in (a) except the retrovirus has been modified further to express on its surface an anticancer cell-binding antibody fragment or an anticancer cell-binding peptide. The virus shown in Figure 4 does not fall within the scope of the claims.

[0115] Figure 5 is a diagrammatic representation of a penton fibre indicating potential fusion sites within the fibre.

[0116] Figure 6 shows fusions between the DNA encoding the Ad5 fibre and an ScFv.

[0117] Figure 7 shows sequences of oligonucleotides used for amplifying the ScFv. All oligonucleotides are presented 5' to 3', the reverse complement of FOR primers are shown and derived amino acid sequences are shown where relevant.

[0118] Figure 8 shows the construction of plasmid pRAS117.

[0119] Figure 9 shows the nucleotide and derived amino acid sequence between the *HindIII* and *EcoRI* sites of pRAS117.

[0120] Figure 10 shows a map of plasmid pRAS117.

[0121] Figure 11 is a diagrammatic representation of the construction of plasmid pRAS118.

[0122] Figure 12 shows the sequences of oligonucleotides for amplifying Ad5 fibre DNA fragments. All oligonucleotides are presented 5' → 3'. The reverse complements of FOR primers are shown. Derived amino acid sequences are shown where relevant.

[0123] Figure 13 shows the nucleotide sequence and deduced amino acid sequence between the *Hind*III site and *Eco*RI site of pRAS111.

[0124] Figure 14 gives a diagrammatic representation of constructing adenovirus carrying a cytotoxic gene.

[0125] Figure 15 gives the nucleotide and amino acid sequences of mouse and humanised HMFG1 variable regions.

Example 1: Fusion sites within the adenovirus Ad5 fibre for binding moieties including single chain Fv (ScFv)

[0126] The Ad5 DNA sequence co-ordinates used here are taken from:

ADRCOMPGE_1: residues 1 to 32760

and ADRCOMPGE_2: residues 32761-35935

[0127] These can be accessed by using program SEQ on the Intelligenetics database.

[0128] The sequence of Ad5 fibre can also be found in Chroboczek, J. and Jacrot, B. (1987) "The sequence of adenovirus fiber: Similarities and differences between serotypes 2 and 5" *Virology* **161**, 549-554 and is available from the EMBL Database, Heidelberg, Germany under accession name ADEFIB.

[0129] Fusion sequences between the shaft and the ScFv are shown in Fig. 6. The fusion sites are at the junctions of the repetitive units of the shaft. Shaft sequences are shown in normal typescript; ScFv sequences are shown in italics. The DNA sequence between the *Pst*I and *Xho*I sites is unique to the ScFv used.

[0130] Fusion A is at the end of the first repetitive unit of the shaft (co-ordinates 31218-9), fusion B at the end of the second (31266-7), fusion C at the third (31323-4), fusion D at the fourth (31368-9), fusion E at the fifth (31413-4), fusion F at the sixth (31458-9), fusion G at the seventh (31503-4), fusion H at the eighth (31551-2), fusion I at the ninth (31596-7), fusion J at the tenth (31641-2), fusion K at the eleventh (31692-3), fusion L at the twelfth (31737-8), fusion M at the thirteenth (31787-8), fusion N at the fourteenth (31836-7), fusion O at the fifteenth (31884-5), fusion P at the sixteenth (31929-30), fusion Q at the seventeenth (31995-6), fusion R at the eighteenth (32040-1), fusion S at the nineteenth (32103-4), fusion T at the twentieth (32151-2), fusion U at the twenty-first (32199-200), and fusion V is at the end of the twenty-second repetitive unit of the shaft (32244-5), the junction between shaft and knob.

Example 2: Preparation of adenovirus expressing an ScFv on its surface

[0131] The genetically modified fibre is introduced into the Ad5 genome by: (a) replacing the fibre gene of plasmid pE4 with the modified fibre by standard recombinant DNA technology and (b) reconstituting the virus by recombination.

[0132] pE4 is a plasmid containing the right hand half of the Ad5 genome, and which has served as the source of the Ad5 fibre gene that we have used. It was provided by Dr Keith Leppard, Biological Sciences, University of Warwick, Coventry, CV4 7AL who has supplied details of its structure. If it is introduced into mammalian cells that contain the remainder of the Ad5 genome, then it is possible to obtain recombinants containing the modification. Most human cell lines can be used for the recombination but HeLa cells are preferred.

[0133] The plasmid pE4 is readily made in the following way. A derivative of pBR322 is made by digesting with *Bst*NI and rejoining using *Xho*I linkers such that the *Bst*NI fragment corresponding to positions 1442-2502 in the pBR322 sequence is removed. DNA from the adenovirus Ad5 strain 309 described by Jones & Shenk (1979) *Cell* **17**, 683-689 is isolated and deproteinated. This DNA is then ligated to *Cla*I linkers and cut with *Eco*RI and *Cla*I. The *Cla*I-*Eco*RI fragment corresponding to the region of 76% of the Ad5 genome to the right hand end is isolated and cloned into the *Eco*RI-*Cla*I sites of the above-mentioned pBR322 derivative to form pE4.

[0134] Adenovirus Type 5 and HeLa cells are available from the American Type Culture Collection, 12301 Packlawn Drive, Rockville, MD 20852-1776, USA under accession numbers ATCC VR-5 and ATCC CCL-2.

Construction of plasmid pRAS117

[0135] Oligonucleotide primers LEADHBACK and LEADbFOR (Figure 7) were used for PCR-mediated amplification of the DNA segment extending from the *Hind*III site of plasmid pRAS111, over the Shine-Dalgarno sequence and the *peB* leader sequence to the *Pst*I site in the ScFv. LEADbFOR directs the incorporation of a *Bgl*II site immediately after the *peB* leader sequence. DNA (100 ng) from plasmid pRAS111 was subjected to 24 rounds of amplification, (94°C, 1 min; 65°C, 1.5 min and 72°C, 2 min) in a 50 µl reaction volume containing 25 pmol of each primer, 250 mM of each dNTP, 67 mM Tris-HCl (pH 8.8), 17 mM (NH₄)₂SO₄, 1.5 mM MgCl₂, 200 mg.ml⁻¹ gelatine and 5 units of *Thermos aquaticus* (Taq) polymerase (Cetus) overlaid with 25 µl paraffin oil. After the reaction, oil was removed by extraction with 500 µl chloroform. The sample was loaded on a 2 % agarose gel, and the amplified fragment was electrophoresed on a piece of NA45 paper (Schleicher and Schuell). Bound DNA was subsequently eluted by immersion in 400 µl 1M NaCl made in TE (10 mM Tris-HCl, pH 7.5, 1 mM EDTA) for 30 min at 70°C. To this was added 800 µl ethanol, and after incubation (2 h, -20°C) the DNA was collected by centrifugation. The pellet was taken up in 50 µl T/E.

[0136] One fifth (10 µl) of the purified amplified fragment was cut with the restriction enzymes *Hind*III and *Pst*I, in a

total volume of 20 µl 50 mM Tris-HCl, pH 7.5, 10 MgCl₂, 100 mM NaCl, 1 mM dithioerythritol containing 10 units of each enzyme. After incubation (1 h, 37°C) the reaction was stopped by incubation at 70°C for 15 minutes.

[0137] The trimmed amplified fragment was cloned between the *Hind*III and *Pst*I sites of pUC8, to generate plasmid pRAS117.

5 [0138] Plasmid pUC8 (1 µg) was cut with *Hind*III and *Pst*I, in a total volume of 20 µl 50 mM Tris-HCl, pH 7.5, 10 MgCl₂, 100 mM NaCl, 1 mM dithioerythritol containing 10 units of each enzyme. After incubation (1 h, 37°C) the reaction was stopped by incubation at 70°C for 15 minutes.

[0139] The ligation reaction contained 1.5 µl of pUC8/*Hind*III, *Pst*I and 3 µl of the amplified leader/*Hind*III, *Pst*I in a total volume of 15 µl containing 70 mM Tris-HCl pH 7.5, 7 mM MgCl₂, 0.7 mM rATP, 4 mM dithioerythritol, 0.5 mg.ml⁻¹ BSA and 10 units of T4 DNA ligase. After incubation (2 h, at room temperature), the reaction was stopped by the addition of 1 µl 500 mM EDTA, pH 8.0 and 14 µl H₂O.

[0140] This ligation mix was used to transform *E. coli*.

[0141] An aliquot (5 µl) of this ligation mix was used to transform a 200 µl aliquot of commercially available competent *E. coli* K12 DH58, 1αF (Life Sciences Inc). After incubation (30 min, 0°C), heat shock (2 min, 42°C), addition of 800 µl L-broth and recovery (37°C, 1 h), cells (100 µl) were spread on L-agar plates containing 100 µg.ml⁻¹ ampicillin containing 50 mM IPTG (isopropyl-β-D-galactopyranoside) and 100 µg.ml⁻¹ X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside). Cells were grown overnight at 37°C, and individual colonies were transferred to fresh L-agar/ampicillin plates. After 6 h growth, colonies were used to inoculate 5 ml aliquots of L-broth containing 100 µg.ml⁻¹ ampicillin. These cells were grown overnight with shaking at 37°C, and used as a source of plasmid DNA.

20 [0142] These cells were used as a source of plasmid DNA.

[0143] Harvested cells were suspended in 360 µl of SET (50 mM sucrose, 10 mM EDTA, 100 mM Tris-HCl, pH 7.5) containing 2 mg.ml⁻¹ hen egg lysozyme, transferred to a 1.5 ml microfuge tube, and diluted by addition of 300 µl 10% Triton X-100. After floating on boiling water for 2 min and cooling for a further minute in ice/water, denatured cell debris was removed by centrifugation (14,000 x g, 20 min) in a microcentrifuge. The majority of the soluble remaining proteins were removed by addition of 300 µl 7.5 M ammonium acetate and centrifugation (14,000 x g, 10 min). Nucleic acids were precipitated by addition of 720 µl cold (-20°C) isopropanol and centrifugation (14,000 x g, 10 min). After rinsing the pellets with ethanol and drying, DNA was solubilised in 60 µl TE containing 170 µg.ml⁻¹ RNase A.

[0144] Restriction enzyme digestions on 5 µl aliquots, using the enzymes *Hind*III and *Bgl*II identified which of these plasmids were pRAS117. The construction scheme is shown in Fig. 8. The nucleotide and derived amino acid sequences between the *Hind*III and *Eco*RI sites of pRAS117 are shown in Fig. 9. A map of plasmid pRAS117 is provided in Fig. 10.

[0145] The nucleotide sequence of the relevant portion of pRAS111, between the *Hind*III site and *Eco*RI, site is given in Figure 13.

35 Construction of plasmid pRAS118 (Figure 11)

[0146] The 130bp *Hind*III-*Pst*I fragment of pRAS117 was used to replace the corresponding fragment of pRAS111, to generate plasmid pRAS118. An aliquot (2 µg) of pRAS111 DNA was cut with *Hind*III and *Pst*I in the conditions used previously, the large fragment was isolated by electrophoresis onto NA45 paper, as described previously, and the DNA was suspended in 10 µl of TE. An aliquot (10 µl) of pRAS117 DNA was cut with *Hind*III and *Pst*I in the conditions used previously, and the small fragment was isolated by electrophoresis onto NA45 paper, as described previously, and the DNA was suspended in 10 µl of TE.

[0147] The isolated pRAS111/*Hind*III-*Pst*I large fragment (1.5 µl) and the isolated pRAS117/*Hind*III-*Pst*I small fragment (3 µl) were mixed and ligated in the conditions previously described.

45 [0148] Transformations, colony handling and DNA preparations were as previously described.

[0149] Restriction enzyme digestions on 5 µl aliquots, using the enzymes *Hind*III, *Pst*I and *Bgl*II identified which of these plasmids were pRAS118. This encodes a NIP-reactive ScFv with a *Bgl*II cloning site immediately downstream of the *peB* leader, suitable for inserting fragments of DNA from Ad5 fibre (and also suitable for fusion of any other desired fusion functions).

50 Amplification of Ad5 fibre DNA fragments

[0150] Fragments of DNA from Ad5 fibre were amplified by PCR using oligonucleotide TAILBACK and oligonucleotide FIBREPFOR, FIBRE3FOR, FIBRE6FOR, FIBRE9FOR, FIBRE12FOR, FIBRE15FOR, FIBRE18FOR, FIBRE21FOR or FIBRE22FOR. Oligonucleotide sequences can be found in Fig. 12.

55 [0151] TAILBACK directs the incorporation of a *Bgl*II site at the base of the fibre, and the FIBREnFOR series primers direct the incorporation of a *Pst*I site at the junctions of repetitive shaft units 3-4 (FIBRE3FOR), 6-7 (FIBRE6FOR), 9-10 (FIBRE9FOR), 12-13 (FIBRE12FOR), 15-16 (FIBRE15FOR), 18-19 (FIBRE18FOR), 21-22 (FIBRE21FOR), be-

tween unit 22 and the knob (FIBRE22FOR) or at the end of the knob sequence (FIBREPFOR).

Fusion of fibre and ScFv

5 [0152] The amplified segments of fibre are trimmed with *Bgl*I and *Pst*I and ligated between the *Bgl*I and *Pst*I sites of plasmid pRAS118. This gives a range of fusions under the transcriptional control of the T7 promoter. Colonies are recovered after transformation of a suitable *E. coli* strain, such as DH5, which does not permit expression of the fusions.

Screening

10 [0153] Colonies containing candidates for fusion are identified by restriction digestion of their plasmid DNAs. These candidate DNAs are used to transform a suitable *E. coli* strain, such as BL21 (DE3), that contains a chromosomal insertion of T7 polymerase under *lac* control. In these cells, induction of expression of T7 polymerase using the gratuitous inducer IPTG causes expression of the fusion proteins. Soluble NIP-reactive material is identified in colonies with correctly assembled fusions. The DNA of these is identified and the NIP-reactive ScFv derived from pRAS111 are replaced with a cell-binding ScFv.

Replacing the fibre:ScFv in plasmid pE4

20 [0154] There is a *Hind*III site approximately half-way along the fibre gene. Fusions with long fibres also contain this *Hind*III site. The fusion is introduced at this site.

Recombination *in vivo* of plasmid pE4-Scfv with the adenovirus genome

25 [0155] To obtain virus particles expressing the ScFv on the penton fibre suitable cells, such as 293 cells, are cotransfected with plasmid pE4-Scfv and plasmid pFG173 as described in Mittal *et al* (1993) *Virus Res.* **28**, 67-90, incorporated herein by reference. Since neither pFG173 nor pE4-ScFv individually is able to generate virus progeny, on transfection of 293 cells viable virus progeny are only produced by *in vivo* recombination between these two plasmids resulting in rescue of the penton fibre-ScFv fusion into the Ad5 genome.

30 [0156] 293 cells are human transformed primary embryonal cells available from the ATCC under accession number ATCC CRL 1573.

[0157] The adenovirus particles made in this way express a NIP-binding ScFv on their surface. Such particles are useful in a two-step targeting approach wherein a target-cell specific binding moiety, such as those identified in Tables 1 and 2, are joined to NIP molecule and targeted to a cell. Once they have localized to the target cell within the patient, the adenovirus displaying NIP-binding ScFv is administered to the patient and binds to the NIP.

Example 3: Insertion of a cytotoxic gene into the E3 region of adenovirus Ad5

40 [0158] In preparation for rescue of the cytotoxic gene into the E3 region of Ad5, the cytotoxic coding sequences were first inserted into a cassette containing the SV40 early promoter and poly A addition sequences to give plasmid pTOX as shown in Figure 14.

[0159] To obtain virus with the cytotoxic gene and SV40 regulatory sequences in the E3 region, 293 cells are cotransfected with plasmids pTOX and pFG173 (Fig 14). The plasmid pFG173 is constructed from pFG140, an infectious plasmid containing the Ad5 d1309 genome in circular form by inserting a *kan*^r gene at the *Eco*RI site as 75.9 m.u. as described in Graham (1984) *EMBO J.* **3**, 2917-2922 and Mitall *et al* (1993) *Virus Res.* **28**, 67-90.

45 [0160] Since neither pFG173 nor pTOX individually is able to generate infectious virus progeny, on transfection of 293 cells viable virus progeny are only produced by *in vivo* recombination between these two plasmids resulting in rescue of the E3 insert into the Ad5 genome.

50 [0161] Viral plaques obtained after cotransfection are isolated and expanded in 293 cells and viral DNA was analyzed on an agarose gel after digestion with *Hind*III. The structure of the desired Ad5-cytotoxic gene recombinant is verified by the presence of diagnostic fragments. One recombinant is plaque purified and used for further study.

Legend to Figure 14

55 [0162] The plasmid pFG173 contains the entire Ad5 genome, except for a 3.2 kb sequence spontaneously deleted between m.u. 75.9-84.9. Plasmids pTOX and pFG173 were used for cotransfection of 293 cells to rescue, by *in vivo* recombination, the cytotoxic gene flanked by SV40 regulatory sequences in the E3 region of Ad5. The resulting Ad5-cytotoxic gene recombinant was named Ad5-TOX. The relative positions of *Hind*III and *Xba*I restriction sites of the

Ad5-TOX genome are shown. The position and orientation of the SV40 promoter, the cytotoxic gene, and the SV40 polyadenylation signal are shown below. Solid bars: luciferase gene; open bars: SV40 promoter and SV40 polyadenylation signal; hatched bars: amp^r and kan^r genes.

[0163] The cytotoxic gene is the cDNA for thymidine kinase.

[0164] Other cytotoxic genes are inserted into the E3 region of Ad5 in an analogous manner.

Example 4: Single chain Fv from the mouse monoclonal antibody HMFG1 and humanised monoclonal antibody Hu HMFG1

[0165] The nucleotide sequences encoding the V_H heavy chains and V_K light chains of HMFG1 and Hu HMFG1 are shown in Figure 15 and are given in Verhoeyen *et al* (1993) *Immunology* **78**, 364-370, incorporated herein by reference.

Legend to Figure 15

[0166] Nucleotide and amino acid sequences of mouse and reshaped HMFG1 variable regions. (a) Heavy chain variable region sequences for mouse and reshaped HMFG1 (Mo V_H-HMFG1 and Hu V_H-HMFG1); (b) mouse and reshaped light chain variable regions respectively (Mo V_K-HMFG1 and Hu V_K-HMFG1). Amino acids numbering and definition of the CDR and framework regions are from Kabat *et al* (1987) *Sequences of Proteins of Immunological Interest*, Edn 4, US Dept of Health and Human Services Public Health Service, NIH, Bethesda, MD 20892, USA.

[0167] The methods described by Bird *et al* (1988) *Science* **242**, 423 or Huston *et al* (1988) *Proc. Natl. Acad. Sci. USA* **85**, 5879 are applied to the nucleotide sequences described in Figure 15 to generate genes encoding ScFv for HMFG and ScFv for Hu HMFG 1. These genes are fused individually into the adenovirus penton fibre gene as described in Examples 1 and 2.

[0168] The amino acid sequences of the V_H and V_L chains of H17E2 are disclosed in "Monoclonal antibodies - applications in clinical oncology", pages 37-43, 1991, A.A. Epenetos, ed., Chapman & Hall, UK.

[0169] Nucleotide sequences encoding the V_H and V_L chains are readily derived from the amino acid sequence using the genetic code and an ScFv can be made from the sequences using the methods of Bird *et al* or Huston *et al* as described above.

Key to Sequence Listing		
Name	SEQ ID No.	
	Nucleotide Sequence	Polypeptide Sequence
Fusion A	1	2
Fusion B	3	4
Fusion C	5	6
Fusion D	7	8
Fusion E	9	10
Fusion F	11	12
Fusion G	13	14
Fusion H	15	16
Fusion I	17	18
Fusion J	19	20
Fusion K	21	22
Fusion L	23	24
Fusion M	25	26
Fusion N	27	28
Fusion O	29	30
Fusion P	31	32
Fusion Q	33	34
Fusion R	35	36
Fusion S	37	38
Fusion T	39	40
Fusion U	41	42
Fusion V	43	44

(continued)

Key to Sequence Listing		
Name	SEQ ID No.	
	Nucleotide Sequence	Polypeptide Sequence
<i>Xho-Eco</i>	45	-
LEADHBACK	46	-
LEADbFOR	47	48
pRAS117	49	50
TAILbBACK	51	52
FIBRE3FOR	53	54
FIBRE6FOR	55	56
FIBRE9FOR	57	58
FIBRE12FOR	59	60
FIBRE15FOR	61	62
FIBRE18FOR	63	64
FIBRE21FOR	65	66
FIBRE22FOR	67	68
FIBREPFOR	69	70
pRAS111	71	72
MoV _H	73	74
MoV _κ	75	76
HuV _H	77	78
HuV _κ	79	80

SEQUENCE LISTING

[0170]

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Imperial Cancer Research Technology Limited

(B) STREET: Sardinia House, Sardinia Street

(C) CITY: London

(E) COUNTRY: United Kingdom

(F) POSTAL CODE (ZIP): WC2A 3NL

(ii) TITLE OF INVENTION: Compounds to target cells

(iii) NUMBER OF SEQUENCES: 80

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

EP 0 672 158 B1

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

10

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Adenovirus
- (B) STRAIN: Ad5

15

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

25

CCT CTA GTT ACC TCC AAT GTG CAG CTG CAG
30
Pro Leu Val Thr Ser Asn Val Gln Leu Gln
1 5 10

30

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: protein

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Pro Leu Val Thr Ser Asn Val Gln Leu Gln

45

1 5 10

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus

(B) STRAIN: Ad5

10 (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..30

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

20 CTC TCT CTG GAC GAG GCC GTC CAG CTG CAG
30
Leu Ser Leu Asp Glu Ala Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 4:

25 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

35

Leu Ser Leu Asp Glu Ala Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 5:

40

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

45

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

50

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

55

(A) ORGANISM: Adenovirus

(B) STRAIN: Ad5

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

10 CCT CTC AAA AAA ACC AAG GTG CAG CTG CAG
30
Pro Leu Lys Lys Thr Lys Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 6:

15

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

25

Pro Leu Lys Lys Thr Lys Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 7:

30

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

45

- (A) ORGANISM: Adenovirus
- (B) STRAIN: Ad5

(ix) FEATURE:

50

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

55

EP 0 672 158 B1

CCC CTC ACA GTT ACC TCA GTG CAG CTG CAG
30
Pro Leu Thr Val Thr Ser Val Gln Leu Gln
1 5 10

5

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

20

Pro Leu Thr Val Thr Ser Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

35

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

40

- (A) ORGANISM: Adenovirus
- (B) STRAIN: Ad5

(ix) FEATURE:

45

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

50

CCT CTA ATG GTC GCG GGC GTG CAG CTG CAG
30
Pro Leu Met Val Ala Gly Val Gln Leu Gln
1 5 10

55

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

EP 0 672 158 B1

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

10 **Pro Leu Met Val Ala Gly Val Gln Leu Gln**
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 11:

15 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

25

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Adenovirus
- (B) STRAIN: Ad5

30

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

40 **CCG CTA ACC GTG CAC GAC GTG CAG CTG CAG**
 30
 Pro Leu Thr Val His Asp Val Gln Leu Gln
 1 5 10

45 (2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: protein

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

EP 0 672 158 B1

Pro Leu Thr Val His Asp Val Gln Leu Gln
1 5 10

5 (2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

20 (vi) ORIGINAL SOURCE:

- (A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

25 (ix) FEATURE:

- (A) NAME/KEY: CDS
(B) LOCATION: 1..30

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

CCC CTC ACA GTG TCA GAA GTG CAG CTG CAG
30
35 Pro Leu Thr Val Ser Glu Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 14:

40 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

50 Pro Leu Thr Val Ser Glu Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 15:

55 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

10

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Adenovirus
- (B) STRAIN: Ad5

15

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..30

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

25

```

      CTC ACC ACC ACC GAT AGC GTG CAG CTG CAG
          30
      Leu Thr Thr Thr Asp Ser Val Gln Leu Gln
          1          5          10

```

(2) INFORMATION FOR SEQ ID NO: 16:

30 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

40

```

      Leu Thr Thr Thr Asp Ser Val Gln Leu Gln
          1          5          10

```

(2) INFORMATION FOR SEQ ID NO: 17:

45

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA (genomic)

55

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus

(B) STRAIN: Ad5

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

CCT CTA ACT ACT GCC ACT GTG CAG CTG CAG
30
Pro Leu Thr Thr Ala Thr Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Pro Leu Thr Thr Ala Thr Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus

(B) STRAIN: Ad5

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

5 CCC ATT TAT ACA CAA AAT GTG CAG CTG CAG
 30
 Pro Ile Tyr Thr Gln Asn Val Gln Leu Gln
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 20:

10

(i) SEQUENCE CHARACTERISTICS:

 (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

20

 Pro Ile Tyr Thr Gln Asn Val Gln Leu Gln
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 21:

25

(i) SEQUENCE CHARACTERISTICS:

 (A) LENGTH: 30 base pairs
30 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

35

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

40

 (A) ORGANISM: Adenovirus
 (B) STRAIN: Ad5

(ix) FEATURE:

45

 (A) NAME/KEY: CDS
 (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

50

 CAT GTA ACA GAC GAC CTA GTG CAG CTG CAG
 30
 His Val Thr Asp Asp Leu Val Gln Leu Gln
 1 5 10

55

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

His Val Thr Asp Asp Leu Val Gln Leu Gln
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Adenovirus
 (B) STRAIN: Ad5

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

GGT GTG ACT ATT AAT AAT GTG CAG CTG CAG
 30
 Gly Val Thr Ile Asn Asn Val Gln Leu Gln
 1 5 10

(2) INFORMATION FOR SEQ ID NO: 24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

Gly Val Thr Ile Asn Asn Val Gln Leu Gln
1 5 10

5 (2) INFORMATION FOR SEQ ID NO: 25:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

20 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

25 (ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

35 GGT TTT GAT TCA CAA GGC GTG CAG CTG CAG
30
Gly Phe Asp Ser Gln Gly Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 26:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

50 Gly Phe Asp Ser Gln Gly Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 27:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

10

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

15

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

25

AGG ATT GAT TCT CAA AAC GTG CAG CTG CAG
30
Arg Ile Asp Ser Gln Asn Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 28:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

40

Arg Ile Asp Ser Gln Asn Val Gln Leu Gln
1 5 10

45

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA (genomic)

55

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

TTT GAT GCT CAA AAC CAA GTG CAG CTG CAG
30
Phe Asp Ala Gln Asn Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

Phe Asp Ala Gln Asn Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

EP 0 672 158 B1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

5 CTT TTT ATA AAC TCA GCC GTG CAG CTG CAG
30
Leu Phe Ile Asn Ser Ala Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 32:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

15

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

20

Leu Phe Ile Asn Ser Ala Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 33:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

30

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

35

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

40

(A) ORGANISM: Adenovirus

(B) STRAIN: Ad5

(ix) FEATURE:

45

(A) NAME/KEY: CDS

(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

50

TCA AAC AAT TCC AAA AAC GTG CAG CTG CAG
30
Ser Asn Asn Ser Lys Asn Val Gln Leu Gln
1 5 10

55

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

Ser Asn Asn Ser Lys Asn Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

(ix) FEATURE:

- (A) NAME/KEY: CDS
(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

GGG TTG ATG TTT GAC GCT GTG CAG CTG CAG
30
Gly Leu Met Phe Asp Ala Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 36:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

Gly Leu Met Phe Asp Ala Val Gln Leu Gln
1 5 10

5 (2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:

10 (A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

20 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

25 (ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

35 CCT AAT GCA CCA AAC ACA GTG CAG CTG CAG
30
Pro Asn Ala Pro Asn Thr Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 38:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

50 Pro Asn Ala Pro Asn Thr Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 39:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

10 (iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

15

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

25

CTA GAA TTT GAT TCA AAC GTG CAG CTG CAG
30
Leu Glu Phe Asp Ser Asn Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 40:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

40

Leu Glu Phe Asp Ser Asn Val Gln Leu Gln
1 5 10

45

(2) INFORMATION FOR SEQ ID NO: 41:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA (genomic)

55

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

CTT AGT TTT GAC AGC ACA GTG CAG CTG CAG
30
Leu Ser Phe Asp Ser Thr Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 42:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

Leu Ser Phe Asp Ser Thr Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 43:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Adenovirus
(B) STRAIN: Ad5

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

5

10

(2) INFORMATION FOR SEQ ID NO: 44:

(i) SEQUENCE CHARACTERISTICS:

15

(ii) MOLECULE TYPE: protein

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

25

(2) INFORMATION FOR SEQ ID NO: 45:

(i) SEQUENCE CHARACTERISTICS:

30

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

40

(vi) ORIGINAL SOURCE:

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

50

(2) INFORMATION FOR SEQ ID NO: 46:

(i) SEQUENCE CHARACTERISTICS:

55

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5 (iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

AGCTAAGCTT GCATGCAAAT TC
22

15 (2) INFORMATION FOR SEQ ID NO: 47:

(i) SEQUENCE CHARACTERISTICS:

20 (A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

30 (ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..27

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

CCA GCG ATG GCC AGA TCT CAG CTG CAG AGCT
31
Pro Ala Met Ala Arg Ser Gln Leu Gln
40 1 5

(2) INFORMATION FOR SEQ ID NO: 48:

45 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

55 Pro Ala Met Ala Arg Ser Gln Leu Gln
1 5

(2) INFORMATION FOR SEQ ID NO: 49:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 132 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 40..132

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

AAGCTTGCAT GCAAATTCTA TTTCAAGGAG ACAGTCATA ATG AAA TAC CTA TTG
 54

Met Lys Tyr Leu Leu
 1 5

CCT ACG GCA GCC GCT GGA TTG TTA TTA CTC GCT GCC CAA CCA GCG ATG
 102

Pro Thr Ala Ala Ala Gly Leu Leu Leu Leu Ala Ala Gln Pro Ala Met
 10 15 20

GCC AGA TCT CAG CTG CAG GTC GAC GGA TCC
 132

Ala Arg Ser Gln Leu Gln Val Asp Gly Ser
 25 30

(2) INFORMATION FOR SEQ ID NO: 50:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

Met Lys Tyr Leu Leu Pro Thr Ala Ala Ala Gly Leu Leu Leu Leu Ala
 1 5 10 15

Ala Gln Pro Ala Met Ala Arg Ser Gln Leu Gln Val Asp Gly Ser
 20 25 30

(2) INFORMATION FOR SEQ ID NO: 51:

(i) SEQUENCE CHARACTERISTICS:

EP 0 672 158 B1

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

10

(iii) ANTI-SENSE: NO

(ix) FEATURE:

15

- (A) NAME/KEY: CDS
- (B) LOCATION: 5..28

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

20

AGCT AGA TCT ATG AAG CGC GCA AGA CCG
28
Arg Ser Met Lys Arg Ala Arg Pro
1 5

25

(2) INFORMATION FOR SEQ ID NO: 52:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

Arg Ser Met Lys Arg Ala Arg Pro
1 5

40

(2) INFORMATION FOR SEQ ID NO: 53:

(i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 41 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

55

(ix) FEATURE:

- (A) NAME/KEY: CDS

(B) LOCATION: 1..33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

5
CCT CTC AAA AAA ACC AAG CAG GTG CAG CTG CAG CAGCCTGG
41
Pro Leu Lys Lys Thr Lys Gln Val Gln Leu Gln
1 5 10

10

(2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:

15
(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

25
Pro Leu Lys Lys Thr Lys Gln Val Gln Leu Gln
1 5 10

25

(2) INFORMATION FOR SEQ ID NO: 55:

(i) SEQUENCE CHARACTERISTICS:

30

(A) LENGTH: 42 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

40

(iii) ANTI-SENSE: NO

(ix) FEATURE:

45
(A) NAME/KEY: CDS
(B) LOCATION: 2..34

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

50
C CCG CTA ACC GTG CAC GAC CAG GTG CAG CTG CAG CAGCCTGG
42
Pro Leu Thr Val His Asp Gln Val Gln Leu Gln
1 5 10

55

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

Pro Leu Thr Val His Asp Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 57:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

CCT CTA ACT ACT GCC ACT CAG GTG CAG CTG CAG CAGCCTGG
41
Pro Leu Thr Thr Ala Thr Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

Pro Leu Thr Thr Ala Thr Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 59:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

10

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..33

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

20

GGT GTG ACT ATT AAT AAT CAG GTG CAG CTG CAG GACCCTGG
41
Gly Val Thr Ile Asn Asn Gln Val Gln Leu Gln
1 5 10

25

(2) INFORMATION FOR SEQ ID NO: 60:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: protein

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

Gly Val Thr Ile Asn Asn Gln Val Gln Leu Gln
1 5 10

40

(2) INFORMATION FOR SEQ ID NO: 61:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

55

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..36

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

5 CCG TTT GAT GCT CAA AAC CAA CAG GTG CAG CTG CAG CAGCC
41
Pro Phe Asp Ala Gln Asn Gln Gln Val Gln Leu Gln
1 5 10

10 (2) INFORMATION FOR SEQ ID NO: 62:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 12 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

Pro Phe Asp Ala Gln Asn Gln Gln Val Gln Leu Gln
1 5 10

25 (2) INFORMATION FOR SEQ ID NO: 63:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 38 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

40 (ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..33

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

50 GGG TTG ATG TTT GAC GCT CAG GTG CAG CTG CAG CAGCC
38
Gly Leu Met Phe Asp Ala Gln Val Gln Leu Gln
1 5 10

55 (2) INFORMATION FOR SEQ ID NO: 64:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

Gly Leu Met Phe Asp Ala Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 40 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 3..35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

GC CTT AGT TTT GAC AGC ACA CAG GTG CAG CTG CAG CAGCC
40
Leu Ser Phe Asp Ser Thr Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

Leu Ser Phe Asp Ser Thr Gln Val Gln Leu Gln
1 5 10

(2) INFORMATION FOR SEQ ID NO: 67:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

5

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

10

(iii) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..45

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

20

GGA AAC AAA AAT AAT GAT AAG CTA ACT TTG CAG GTG CAG CTG CAG
 45
 Gly Asn Lys Asn Asn Asp Lys Leu Thr Leu Gln Val Gln Leu Gln
 1 5 10 15

CAGCC
 50

25

(2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

40

Gly Asn Lys Asn Asn Asp Lys Leu Thr Leu Gln Val Gln Leu Gln
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO: 69:

45

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 43 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

55

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 3..17

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

5

CA TAC ATT GCC CAA GAA TAACAGGTGC AGCTGCAGCA GCCTGG
43
Tyr Ile Ala Gln Glu
1 5

10

(2) INFORMATION FOR SEQ ID NO: 70:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

25

Tyr Ile Ala Gln Glu
1 5

(2) INFORMATION FOR SEQ ID NO: 71:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 858 base pairs

(B) TYPE: nucleic acid (C STRANDEDNESS: double

(D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

40

(iii) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

45

(B) LOCATION: 40..846

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

50

55

AAGCTTGCAT GCAAATTCTA TTTCAGGAG ACAGTCATA ATG AAA TAC CTA TTG
 54
 Met Lys Tyr Leu Leu
 1 5

5
 CCT ACG GCA GCC GCT GGA TTG TTA TTA CTC GCT GCC CAA CCA GCG ATG
 102
 Pro Thr Ala Ala Ala Gly Leu Leu Leu Leu Ala Ala Gln Pro Ala Met
 10 15 20

10
 GCC CAG GTG CAG CTG CAG CAG CCT GGG GCT GAG CTT GTG AAG CCT GGG
 150
 Ala Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly
 25 30 35

15
 GCT TCA GTG AAG CTG TCC TGC AAG GCT TCT GGC TAC ACC TTC ACC AGC
 198
 Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser
 40 45 50

20
 TAC TGG ATG CAC TGG GTG AAG CAG AGG CCT GGA CGA GGC CTT GAG TGG
 246
 Tyr Trp Met His Trp Val Lys Gln Arg Pro Gly Arg Gly Leu Glu Trp
 55 60 65

25
 ATT GGA AGG ATT GAT CCT AAT AGT GGT GGT ACT AAG TAC AAT GAG AAG
 294
 Ile Gly Arg Ile Asp Pro Asn Ser Gly Gly Thr Lys Tyr Asn Glu Lys
 70 75 80 85

30
 TTC AAG AGC AAG GCC ACA CTG ACT GTA GAC AAA CCC TCC AGC ACA GCC
 342
 Phe Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Pro Ser Ser Thr Ala
 90 95 100

35
 TAC ATG CAG CTC AGC AGC CTG ACA TCT GAG GAC TCT GCG GTC TAT TAT
 390
 Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr
 105 110 115

40
 TGT GCA AGA TAC GAT TAC TAC GGT AGT AGC TAC TTT GAC TAC TGG GGC
 438
 Cys Ala Arg Tyr Asp Tyr Tyr Gly Ser Ser Tyr Phe Asp Tyr Trp Gly
 120 125 130

45
 CAA GGG ACC ACG GTC ACC GTC TCC TCA GGT GGA GGC GGT TCA GGC GGA
 486
 Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly
 135 140 145

50
 GGT GGC TCT GGC GGT GGC GGA TCC CAG GCT GTT GTG ACT CAG GAA TCT
 534
 Gly Gly Ser Gly Gly Gly Gly Ser Gln Ala Val Val Thr Gln Glu Ser
 150 155 160 165

55
 GCA CTC ACC ACA TCA CCT GGT GAA ACA GTC ACA CTC ACT TGT CGC TCA
 582
 Ala Leu Thr Thr Ser Pro Gly Glu Thr Val Thr Leu Thr Cys Arg Ser

170 175 180
 AGT ACT GGG GCT GTT ACA ACT AGT AAC TAT GCC AAC TGG GTC CAA GAA
 630
 5 Ser Thr Gly Ala Val Thr Thr Ser Asn Tyr Ala Asn Trp Val Gln Glu
 185 190 195
 AAA CCA GAT CAT TTA TTC ACT GGT CTA ATA GGT GGT ACC AAC AAC CGA
 678
 10 Lys Pro Asp His Leu Phe Thr Gly Leu Ile Gly Gly Thr Asn Asn Arg
 200 205 210
 GCT CCA GGT GTT CCT GCC AGA TTC TCA GGC TCC CTG ATT GGA GAC AAG
 726
 15 Ala Pro Gly Val Pro Ala Arg Phe Ser Gly Ser Leu Ile Gly Asp Lys
 215 220 225
 GCT GCC CTC ACC ATC ACA GGG GCA CAG ACT GAG CAT GAG GCA ATA TAT
 774
 20 Ala Ala Leu Thr Ile Thr Gly Ala Gln Thr Glu Asp Glu Ala Ile Tyr
 230 235 240 245
 TTC TGT GCT CTA TGG TAC ACC AAC CAC TGG GTG TTC GGT GGA GGA ACC
 822
 25 Phe Cys Ala Leu Trp Tyr Ser Asn His Trp Val Phe Gly Gly Gly Thr
 250 255 260
 AAA CTG ACT GTC CTA GGT CTC GAG TAATAAGAAT TC
 858
 Lys Leu Thr Val Leu Gly Leu Glu
 265

(2) INFORMATION FOR SEQ ID NO: 72:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 269 amino acids

(B) TYPE: amino acid

35

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

40

45

50

55

EP 0 672 158 B1

Met Lys Tyr Leu Leu Pro Thr Ala Ala Ala Gly Leu Leu Leu Leu Ala
1 5 10 15
Ala Gln Pro Ala Met Ala Gln Val Gln Leu Gln Gln Pro Gly Ala Glu
5 20 25 30
Leu Val Lys Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly
35 40 45
Tyr Thr Phe Thr Ser Tyr Trp Met His Trp Val Lys Gln Arg Pro Gly
10 50 55 60
Arg Gly Leu Glu Trp Ile Gly Arg Ile Asp Pro Asn Ser Gly Gly Thr
65 70 75 80
Lys Tyr Asn Glu Lys Phe Lys Ser Lys Ala Thr Leu Thr Val Asp Lys
15 85 90 95
Pro Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp
100 105 110
Ser Ala Val Tyr Tyr Cys Ala Arg Tyr Asp Tyr Tyr Gly Ser Ser Tyr
20
115 120 125
Phe Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly
25 130 135 140
Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gln Ala Val
145 150 155 160
Val Thr Gln Glu Ser Ala Leu Thr Thr Ser Pro Gly Glu Thr Val Thr
30 165 170 175
Leu Thr Cys Arg Ser Ser Thr Gly Ala Val Thr Thr Ser Asn Tyr Ala
180 185 190
Asn Trp Val Gln Glu Lys Pro Asp His Leu Phe Thr Gly Leu Ile Gly
35 195 200 205
Gly Thr Asn Asn Arg Ala Pro Gly Val Pro Ala Arg Phe Ser Gly Ser
210 215 220
Leu Ile Gly Asp Lys Ala Ala Leu Thr Ile Thr Gly Ala Gln Thr Glu
40 225 230 235 240
Asp Glu Ala Ile Tyr Phe Cys Ala Leu Trp Tyr Ser Asn His Trp Val
245 250 255
Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Leu Glu
45 260 265

(2) INFORMATION FOR SEQ ID NO: 73:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 354 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..354

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

```

15      CAG GTT CAG CTG CAG CAG TCT GGA GCT GAG CTG ATG AAG CCT GGG GCC
          48
      Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Met Lys Pro Gly Ala
          1           5           10           15

20      TCA GTG AAG ATA TCC TGC AAG GCT ACT GGC TAC ACA TTC AGT GCC TAC
          96
      Ser Val Lys Ile Ser Cys Lys Ala Thr Gly Tyr Thr Phe Ser Ala Tyr
          20           25           30

25      TGG ATA GAG TGG GTA AAG CAG AGG CCT GGA CAT GGC CTT GAG TGG ATT
          144
      Trp Ile Glu Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp Ile

          35           40           45

30      GGA GAG ATT TTA CCT GGA AGT AAT AAT TCT AGA TAC AAT GAG AAG TTC
          192
      Gly Glu Ile Leu Pro Gly Ser Asn Asn Ser Arg Tyr Asn Glu Lys Phe
          50           55           60

35      AAG GGC AAG GCC ACA TTC ACT GCT GAT ACA TCC TCC AAC ACA GCC TAC
          240
      Lys Gly Lys Ala Thr Phe Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr
          65           70           75           80

40      ATG CAA CTC AGC AGC CTG ACA TCT GAG GAC TCT GCC GTC TAT TAC TGT
          288
      Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
          85           90           95

45      TCA AGG TCC TAC GAC TTT GCC TGG TTT GCT TAC TGG GGC CAA GGG ACT
          336
      Ser Arg Ser Tyr Asp Phe Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr
          100           105           110

50      CCG GTC ACT GTC TCT GCA
          354
      Pro Val Thr Val Ser Ala
          115

```

(2) INFORMATION FOR SEQ ID NO: 74:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 118 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:

5 Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Met Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Ile Ser Cys Lys Ala Thr Gly Tyr Thr Phe Ser Ala Tyr
 20 25 30
 10 Trp Ile Glu Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile Leu Pro Gly Ser Asn Asn Ser Arg Tyr Asn Glu Lys Phe
 50 55 60
 15 Lys Gly Lys Ala Thr Phe Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 20 Ser Arg Ser Tyr Asp Phe Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr
 100 105 110
 Pro Val Thr Val Ser Ala
 115

25 (2) INFORMATION FOR SEQ ID NO: 75:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 342 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

40 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(ix) FEATURE:

45 (A) NAME/KEY: CDS
 (B) LOCATION: 1..342

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

50

55

5 GAC ATT GTG ATG TCA CAG TCT CCA TCC TCC CTA GCT GTG TCA GTT GGA
 48
 Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
 1 5 10 15
 10 GAG AAG GTT ACT ATG AGC TGC AAG TCC AGT CAG AGC CTT TTA TAT AGT
 96
 Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 15 AGC AAT CAA AAG ATC TAC TTG GCC TGG TAC CAG CAG AAA CCA GGG CAG
 144
 Ser Asn Gln Lys Ile Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 20 TCT CCT AAA CTG CTG ATT TAC TGG GCA TCC ACT AGG GAA TCT GGG GTC
 192
 Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 25 CCT GAT CGC TTC ACA GGC GGT GGA TCT GGG ACA GAT TTC ACT CTC ACC
 240
 Pro Asp Arg Phe Thr Gly Gly Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 30 ATC AGC AGT GTG AAG GCT GAA GAC CTG GCA GTT TAT TAC TGT CAG CAA
 288
 Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 TAT TAT AGA TAT CCT CGG ACG TTC GGT GGA GGC ACC AAG CTG GAA ATC
 336
 Tyr Tyr Arg Tyr Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
 100 105 110
 AAA CGG
 342
 Lys Arg

35 (2) INFORMATION FOR SEQ ID NO: 76:

 (i) SEQUENCE CHARACTERISTICS:

40 (A) LENGTH: 114 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

 (ii) MOLECULE TYPE: protein

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

50

55

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
 1 5 10 15
 5 Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30
 Ser Asn Gln Lys Ile Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 10 Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 Pro Asp Arg Phe Thr Gly Gly Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 15 Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
 85 90 95
 Tyr Tyr Arg Tyr Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
 100 105 110
 20 Lys Arg

(2) INFORMATION FOR SEQ ID NO: 77:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 354 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..354

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

45 CAG GTG CAG CTG GTG CAG TCT GGG GCA GAG GTG AAA AAG CCT GGG GCC
 48
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 50 TCA GTG AAG GTG TCC TGC AAG GCT TCT GGC TAC ACC TTC AGT GCC TAC
 96
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Ser Ala Tyr
 20 25 30
 TGG ATA GAG TGG GTG CGC CAG GCT CCA GGA AAG GGC CTC GAG TGG GTC
 144
 55 Trp Ile Glu Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

	35					40					45					
	GGA	GAG	ATT	TTA	CCT	GGA	AGT	AAT	AAT	TCT	AGA	TAC	AAT	GAG	AAG	TTC
5	Gly	Glu	Ile	Leu	Pro	Gly	Ser	Asn	Asn	Ser	Arg	Tyr	Asn	Glu	Lys	Phe
	192	50				55					60					
	AAG	GGC	CGA	GTG	ACA	GTC	ACT	AGA	GAC	ACA	TCC	ACA	AAC	ACA	GCC	TAC
10	Lys	Gly	Arg	Val	Thr	Val	Thr	Arg	Asp	Thr	Ser	Thr	Asn	Thr	Ala	Tyr
	65				70						75				80	
	ATG	GAG	CTC	AGC	AGC	CTG	AGG	TCT	GAG	GAC	ACA	GCC	GTC	TAT	TAC	TGT
	288															
15	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
					85					90					95	
	GCA	AGA	TCC	TAC	GAC	TTT	GCC	TGG	TTT	GCT	TAC	TGG	GGC	CAA	GGG	ACT
	336															
	Ala	Arg	Ser	Tyr	Asp	Phe	Ala	Trp	Phe	Ala	Tyr	Trp	Gly	Gln	Gly	Thr
				100					105					110		
20	CTG	GTC	ACA	GTC	TCC	TCA										
	354															
	Leu	Val	Thr	Val	Ser	Ser										
				115												

25 (2) INFORMATION FOR SEQ ID NO: 78:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 118 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

[illegible]

(2) INFORMATION FOR SEQ ID NO: 79:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 342 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..342

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

```

GAC ATC CAG ATG ACC CAG AGC CCA AGC AGC CTG AGC GCC AGC GTG GGT
 48
Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1      5      10      15
GAC AGA GTG ACC ATC ACC TGT AAG TCC AGT CAG AGC CTT TTA TAT AGT
 96
Asp Arg Val Thr Ile Thr Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20      25      30
AGC AAT CAA AAG ATC TAC TTG GCC TGG TAC CAG CAG AAG CCA GGT AAG
144
Ser Asn Gln Lys Ile Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
 35      40      45
GCT CCA AAG CTG CTG ATC TAC TGG GCA TCC ACT AGG GAA TCT GGT GTG
192
Ala Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50      55      60
CCA AGC AGA TTC AGC GGT AGC GGT AGC GGT ACC GAC TTC ACC TTC ACC
240
Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr
 65      70      75      80
ATC AGC AGC CTC CAG CCA GAG GAC ATC GCC ACC TAC TAC TGC CAG CAA
288
Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln
 85      90      95
TAT TAT AGA TAT CCT CGG ACG TTC GGC CAA GGG ACC AAG GTG GAA ATC
336
Tyr Tyr Arg Tyr Pro Arg Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
100      105      110
AAA CGT
342
Lys Arg

```

(2) INFORMATION FOR SEQ ID NO: 80:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 114 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

```

10      Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
        1           5           10           15
      Asp Arg Val Thr Ile Thr Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
        20           25           30
15      Ser Asn Gln Lys Ile Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys
        35           40           45
      Ala Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
        50           55           60
20      Pro Ser Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr
        65           70           75           80
      Ile Ser Ser Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln
        85           90           95
25      Tyr Tyr Arg Tyr Pro Arg Thr Phe Gly Gln Gly Thr Lys Val Glu Ile
        100          105          110
      Lys Arg
  
```

30

Claims

1. An adenovirus, or adenovirus-like particle, or a replication defective derivative thereof comprising a modified binding specificity conferred by a binding moiety allowing the adenovirus or adenovirus-like particle to bind to a target cell, wherein the binding moiety is a peptide and is fused to the penton fibre of the adenovirus or adenovirus-like particle.
2. An adenovirus or adenovirus-like particle according to Claim 1 wherein the binding moiety is a monoclonal antibody, an ScFv, a dAb, or a minimal recognition unit of an antibody.
3. An adenovirus or adenovirus-like particle according to Claim 1 wherein the binding moiety is at least part of a ligand of a target cell-specific cell-surface receptor.
4. An adenovirus or adenovirus-like particle according to Claim 2 or 3 wherein the binding moiety recognises a target cell-specific surface antigen.
5. An adenovirus or adenovirus-like particle according to Claim 3 wherein the target cell-specific cell-surface receptor is any one of GnRH receptor, MSH receptor and somatostatin receptor.
6. An adenovirus or adenovirus-like particle according to any one of Claims 1 to 5 containing a gene suitable for gene therapy.
7. An adenovirus or adenovirus-like particle according to Claim 6 wherein the gene encodes a molecule having a directly or indirectly cytotoxic function.
8. An adenovirus or adenovirus-like particle according to Claim 7 wherein the gene encodes any one of interleukin-2, tumour necrosis factor, interferon-gamma, ribonuclease and deoxyribonuclease.

9. An adenovirus or adenovirus-like particle according to Claim 7 wherein the gene encodes an enzyme capable of converting a relatively non-toxic pro-drug into a cytotoxic drug.
- 5 10. An adenovirus or adenovirus-like particle according to Claim 9 wherein the gene is either cytosine deaminase or thymidine kinase.
11. An adenovirus or adenovirus-like particle according to Claim 6 wherein the gene overcomes a defect in a gene in the target cell.
- 10 12. An adenovirus or adenovirus-like particle according to Claim 11 wherein the gene is any one of CFTR, dystrophin and haemoglobin A.
13. An adenovirus, or adenovirus-like particle, according to any one of Claims 1 to 19 further containing nucleic acid wherein the said adenovirus or adenovirus-like particle is adapted to deliver the said nucleic acid to the target cell.
- 15 14. An adenovirus or adenovirus-like particle according to Claim 1 wherein the binding moiety is fused to the penton fibre protein at any one or more of the junctions of the repetitive units of the shaft.
15. An adenovirus according to Claim 14 wherein the binding moiety is a ScFv.
- 20 16. An adenovirus according to Claim 15 wherein the ScFv binds to a tumour cell antigen.
17. An adenovirus or adenovirus-like particle according to any one of Claims 1 to 16 wherein the binding moiety is a polypeptide.
- 25 18. An adenovirus or adenovirus-like particle according to any one of Claims 1 to 17 for use in medicine.
19. A nucleotide sequence encoding the fusion of the binding moiety and the penton fibre said fusion being that present on the adenovirus or adenovirus-like particle as defined in any one of Claims 1 to 17.
- 30 20. A nucleotide sequence defined in Claim 19 additionally comprising the remainder of the genome of the adenovirus or adenovirus-like particle.
21. A nucleotide sequence encoding a virus or virus-like particle according to any one of Claims 1 to 17.
- 35 22. A system of components for use in a therapeutic method comprising an adenovirus or adenovirus-like particle according to Claim 14 or 15 and a pro-drug.
23. A method for producing an adenovirus or adenovirus-like particle according to any of Claims 1 to 17 in cell culture, the method comprising (1) genetically modifying the adenovirus or adenovirus-like particle to produce a binding moiety, (2) infecting cells with the genetically modified adenovirus or adenovirus-like particle, (3) culturing the cells until the adenovirus or adenovirus-like particle reaches a sufficiently high titre and (4) harvesting and substantially purifying the genetically modified adenovirus or adenovirus-like particle.
- 40 24. A pharmaceutical composition comprising an adenovirus or adenovirus-like particle according to any one of Claims 1 to 17 and a pharmaceutical carrier.
25. Use of an adenovirus or adenovirus-like particle according to Claim 7 in the manufacture of a medicament for destroying target cells in a mammal.
- 50 26. Use of an adenovirus or adenovirus-like particle according to Claim 9 or 10 in the manufacture of a medicament for destroying target cells in a mammal wherein said mammal will be administered said pro-drug.
27. Use of an adenovirus or adenovirus-like particle according to Claim 11 or 12 in the manufacture of a medicament for treating a mammal having a defective gene.
- 55

Patentansprüche

1. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen oder ein replikationsdefektes Derivat davon, das eine modifizierte Bindungsspezifität aufweist, die von einer Bindungseinheit verliehen wird, die es dem Adenovirus oder dem Adenovirus-ähnlichen Teilchen ermöglicht, an eine Zielzelle zu binden, wobei die Bindungseinheit ein Peptid ist und an die Penton-Faser des Adenovirus oder des Adenovirus-ähnlichen Teilchens fusioniert ist.
2. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 1, wobei die Bindungseinheit ein monoklonaler Antikörper, ein ScFv, ein dAb oder eine minimale Erkennungseinheit eines Antikörpers ist.
3. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 1, wobei die Bindungseinheit wenigstens ein Teil eines Liganden eines zielzellenspezifischen Zelloberflächen-Rezeptors ist.
4. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 2 oder 3, wobei die Bindungseinheit ein zielzellenspezifisches Oberflächenantigen erkennt.
5. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 3, wobei der zielzellenspezifische Zelloberflächen-Rezeptor einer von dem GnRH-Rezeptor, MSH-Rezeptor und Somatostatin-Rezeptor ist.
6. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach irgendeinem der Ansprüche 1 bis 5, das ein Gen enthält, das für die Gentherapie geeignet ist.
7. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 6, wobei das Gen für ein Molekül mit einer direkten oder indirekten cytotoxischen Funktion codiert.
8. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 7, wobei das Gen für eines von Interleukin-2, Tumornekrosefaktor, Interferon-Gamma, Ribonuclease und Desoxyribonuclease codiert.
9. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 7, wobei das Gen für ein Enzym codiert, das in der Lage ist, eine relativ nicht-toxische Vorstufe in eine cytotoxische Droge zu überführen.
10. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 9, wobei das Gen entweder Cytosindeaminase oder Thymidinkinase ist.
11. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 6, wobei das Gen einen Defekt in einem Gen in der Zielzelle beseitigt.
12. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 11, wobei das Gen irgendeines von CFTR, Dystrophin und Hämoglobin A ist.
13. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach irgendeinem der Ansprüche 1 bis 12, das außerdem eine Nukleinsäure enthält, wobei das Adenovirus oder das Adenovirus-ähnliche Teilchen geeignet ist, die Nukleinsäure in die Zielzelle einzuführen.
14. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 1, wobei die Bindungseinheit an das Penton-Faser-Protein an irgendeiner oder mehreren der Verknüpfungen der sich wiederholenden Einheiten des Schafte fusioniert ist.
15. Ein Adenovirus nach Anspruch 14, wobei die Bindungseinheit ein ScFv ist.
16. Ein Adenovirus gemäß Anspruch 15, wobei das ScFv an ein Tumorzell-Antigen bindet.
17. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach irgendeinem der Ansprüche 1 bis 16, wobei die Bindungseinheit ein Polypeptid ist.
18. Ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach irgendeinem der Ansprüche 1 bis 17 für die Verwendung in der Medizin.

19. Eine Nucleotidsequenz, die für die Fusion der Bindungseinheit mit der Penton-Faser codiert, wobei die Fusion so an dem Adenovirus oder an dem Adenovirus-ähnlichen Teilchen vorliegt, wie in irgendeinem der Ansprüche 1 bis 17 definiert ist.
- 5 20. Eine Nucleotidsequenz nach Anspruch 19, die zusätzlich den Rest des Genoms des Adenovirus oder des Adenovirus-ähnlichen Teilchens umfaßt.
21. Eine Nucleotidsequenz, die für ein Virus oder für ein Virus-ähnliches Teilchen nach irgendeinem der Ansprüche 1 bis 17 codiert.
- 10 22. Ein System aus Komponenten für die Verwendung bei einem therapeutischen Verfahren, das ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach Anspruch 14 oder 15 und eine Vordroge umfaßt.
23. Verfahren zur Herstellung eines Adenovirus oder eines Adenovirus-ähnlichen Teilchens nach irgendeinem der Ansprüche 1 bis 17 in einer Zellkultur, wobei das Verfahren umfaßt (1) die genetische Modifizierung des Adenovirus oder des Adenovirus-ähnlichen Teilchens, um eine Bindungseinheit zu erzeugen, (2) das Infizieren der Zellen mit dem genetisch modifizierten Adenovirus oder dem Adenovirus-ähnlichen Teilchen, (3) Züchten der Zellen, bis das Adenovirus oder das Adenovirus-ähnliche Teilchen einen ausreichend hohen Titer erreicht und (4) Ernten und im wesentlichen Reinigen des genetisch veränderten Adenovirus oder des Adenovirus-ähnlichen Teilchens.
- 15 24. Eine pharmazeutische Zusammensetzung, die ein Adenovirus oder ein Adenovirus-ähnliches Teilchen nach irgendeinem der Ansprüche 1 bis 17 und einen pharmazeutischen Träger umfaßt.
- 25 25. Verwendung eines Adenovirus oder eines Adenovirus-ähnlichen Teilchens nach Anspruch 7 bei der Herstellung eines Medikaments zur Zerstörung von Zielzellen in einem Säugetier.
26. Verwendung eines Adenovirus oder eines Adenovirus-ähnlichen Teilchens nach Anspruch 9 oder 10 bei der Herstellung eines Medikaments zur Zerstörung von Zielzellen in einem Säugetier, wobei dem Säugetier die Vordroge verabreicht wird.
- 30 27. Verwendung eines Adenovirus oder eines Adenovirus-ähnlichen Teilchens nach Anspruch 11 oder 12 bei der Herstellung eines Medikaments zur Behandlung eines Säugetiers mit einem defekten Gen.

35 **Revendications**

1. Adénovirus, ou particule de type adénovirus, ou dérivé défectueux dans la réplication de ceux-ci, comprenant une spécificité de liaison modifiée conférée par un élément de liaison permettant à l'adénovirus ou à la particule de type adénovirus de se lier à une cellule cible, dans lequel l'élément de liaison est un peptide et est fusionné à la fibre du penton de l'adénovirus ou de la particule de type adénovirus.
- 40 2. Adénovirus ou particule de type adénovirus selon la revendication 1, dans lequel l'élément de liaison est un anticorps monoclonal, un ScFv, un dAb, ou une unité de reconnaissance minimale d'un anticorps.
- 45 3. Adénovirus ou particule de type adénovirus selon la revendication 1, dans lequel l'élément de liaison est au moins une partie d'un ligand d'un récepteur de surface cellulaire spécifique d'une cellule cible.
4. Adénovirus ou particule de type adénovirus selon la revendication 2 ou la revendication 3, dans lequel l'élément de liaison reconnaît un antigène de surface spécifique d'une cellule cible.
- 50 5. Adénovirus ou particule de type adénovirus selon la revendication 3, dans lequel le récepteur de surface cellulaire spécifique d'une cellule cible est l'un quelconque parmi le récepteur de la GnRH, le récepteur de la MSH et le récepteur de la somatostatine.
- 55 6. Adénovirus ou particule de type adénovirus selon l'une quelconque des revendications 1 à 5, contenant un gène convenant à une thérapie génique.
7. Adénovirus ou particule de type adénovirus selon la revendication 6, dans lequel le gène code pour une molécule

ayant une fonction directement ou indirectement cytotoxique.

- 5 8. Adénovirus ou particule de type adénovirus selon la revendication 7, dans lequel le gène code pour l'un quelconque parmi l'interleukine-2, le facteur nécrosant des tumeurs, l'interféron gamma, la ribonucléase et la désoxyribonucléase.
9. Adénovirus ou particule de type adénovirus selon la revendication 7, dans lequel le gène code pour une enzyme capable de convertir un promédicament relativement non toxique en un médicament cytotoxique.
- 10 10. Adénovirus ou particule de type adénovirus selon la revendication 9, dans lequel le gène est soit la cytosine-désaminase, soit la thymidine-kinase.
11. Adénovirus ou particule de type adénovirus selon la revendication 6, dans lequel le gène vient à bout d'un défaut dans un gène dans la cellule cible.
- 15 12. Adénovirus ou particule de type adénovirus selon la revendication 11, dans lequel le gène est l'un quelconque parmi le CFTR, la dystrophine et l'hémoglobine A.
- 20 13. Adénovirus ou particule de type adénovirus selon l'une quelconque des revendications 1 à 19, contenant en plus de l'acide nucléique, dans lequel ledit adénovirus ou particule de type adénovirus est adapté à la délivrance dudit acide nucléique à la cellule cible.
- 25 14. Adénovirus ou particule de type adénovirus selon la revendication 1, dans lequel l'élément, de liaison est fusionné à la protéine de fibre du penton au niveau de l'une quelconque ou de plusieurs jonctions des unités répétitives de l'axe.
15. Adénovirus selon la revendication 14, dans lequel l'élément de liaison est un ScFv.
- 30 16. Adénovirus selon la revendication 15, dans lequel le ScFv se lie à un antigène de cellule tumorale.
17. Adénovirus ou particule de type adénovirus selon l'une quelconque des revendications 1 à 16, dans lequel l'élément de liaison est un polypeptide.
- 35 18. Adénovirus ou particule de type adénovirus selon l'une quelconque des revendications 1 à 17 destiné à une utilisation en médecine.
- 40 19. Séquence de nucléotides encodant la fusion de l'élément de liaison et de la fibre du penton, ladite fusion étant celle présente sur l'adénovirus ou la particule de type adénovirus tel que défini dans l'une quelconque des revendications 1 à 17.
20. Séquence de nucléotides définie dans la revendication 19, comprenant en plus le reste du génome de l'adénovirus ou de la particule de type adénovirus.
- 45 21. Séquence de nucléotides encodant un virus ou une particule de type virus selon l'une quelconque des revendications 1 à 17.
22. Système de composants pour une utilisation dans un procédé thérapeutique comprenant un adénovirus ou une particule de type adénovirus selon la revendication 14 ou la revendication 15 et un promédicament.
- 50 23. Procédé pour produire un adénovirus ou une particule de type adénovirus selon l'une quelconque des revendications 1 à 17 en culture cellulaire, le procédé comprenant les étapes consistant à (1) modifier génétiquement l'adénovirus ou la particule de type adénovirus pour produire un élément de liaison, (2) infecter des cellules avec l'adénovirus ou la particule de type adénovirus génétiquement modifié, (3) mettre les cellules en culture jusqu'à ce que l'adénovirus ou la particule de type adénovirus atteigne un titre suffisamment élevé et (4) récolter et sensiblement purifier l'adénovirus ou la particule de type adénovirus génétiquement modifié.
- 55 24. Composition pharmaceutique comprenant un adénovirus ou une particule de type adénovirus selon l'une quelconque des revendications 1 à 17 et un véhicule pharmaceutique.

25. Utilisation d'un adénovirus ou d'une particule de type adénovirus selon la revendication 7 dans la fabrication d'un médicament destiné à détruire des cellules cibles chez un mammifère.

26. Utilisation d'un adénovirus ou d'une particule de type adénovirus selon la revendication 9 ou la revendication 10 dans la fabrication d'un médicament destiné à détruire des cellules cibles chez un mammifère, dans laquelle ledit mammifère se verra administrer ledit promédicament.

27. Utilisation d'un adénovirus ou d'une particule de type adénovirus selon la revendication 11 ou la revendication 12 dans la fabrication d'un médicament destiné à traiter un mammifère ayant un gène défectueux.




Fig. 1(a)




Fig. 1(b)




Fig. 1(c)




Fig. 2(a)




Fig. 2(b)




Fig. 2(c)




Fig. 3(a)




Fig. 3(b)




Fig. 4(a)




Fig. 4(b)




Figure 5

Fusion A

		1	2	3	4		108	109					
P	L	V	T	S	N	V	Q	L	QL	E	*	*
CCTCTAGTTACCTCCAATGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC													
31210						PstI ..		.XhoI		EcoRI			

Fusion B

				1	2	3	4		108	109			
L	S	L	D	E	A	V	Q	L	QL	E	*	*
CTCTCTCTGGACGAGGCCGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC													
31260						PstI ..		.XhoI		EcoRI			

Fusion C

				1	2	3	4		108	109			
P	L	K	K	T	K	V	Q	L	QL	E	*	*
CCTCTCAAAAAACCAAGGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC													
31310	31320					PstI ..		.XhoI		EcoRI			

Fusion D

				1	2	3	4		108	109			
P	L	T	V	T	S	V	Q	L	QL	E	*	*
CCCCTCACAGTTACCTCAGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC													
31360						PstI ..		.XhoI		EcoRI			

Fusion E

				1	2	3	4		108	109			
P	L	M	V	A	D	V	Q	L	QL	E	*	*
CCTCTAATGGTCGCGGGCGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC													
31400	31410					PstI ..		.XhoI		EcoRI			

Figure 6 (Page 1 of 5)

Fusion F

1 2 3 4 108 109
 P L T V H D V Q L Q L E * *
 CCGCTAACCGTGACGACGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC
 31450 | PstI .. .XhoI EcoRI

Fusion G

1 2 3 4 108 109
P L T V S E V Q L Q L E * *
CCCCTCACAGTGTCTCAGAAAGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC
31490 31500 | PstI .. .XhoI EcoRI

Fusion H

1 2 3 4 108 109
 L T T T D S V Q L Q L E * *
 CTCACCACCACCGATAGCGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC
 31540 31550 | PstI .. XhoI EcoRI

Fusion I

1 2 3 4 108 109
 P L T T A T V Q L Q L E * *
 CCTCTAACTACTGCCACTGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC
 31590 | PstI .. .XhoI EcoRI

Fusion J

1 2 3 4 108 109
P I Y T Q N V Q L QL E * *
CCCATTTTATACACAAAATGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC
31630 31640 | PstI .. XhoI EcoRI

Figure 6 (Page 2 of 5)

Fusion K

	1	2	3	4		108	109
H	V	T	D	D	L	V	Q
					L	QL
							E * *
CATGTAACAGACGACCTAGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
31730				PstI ..		.XhoI	EcoRI

Fusion L

	1	2	3	4		108	109
G	V	T	I	N	N	V	Q
					L	QL
							E * *
GGTGTGACTATTAATAATGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
31730				PstI ..		.XhoI	EcoRI

Fusion M

	1	2	3	4		108	109
G	F	D	S	Q	G	V	Q
					L	QL
							E * *
GGTTTTGATTTCACAAGGCGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
31780				PstI ..		.XhoI	EcoRI

Fusion N

	1	2	3	4		108	109
R	I	D	S	Q	N	V	Q
					L	QL
							E * *
AGGATTGATTCTCAAACGTCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
31830				PstI ..		.XhoI	EcoRI

Fusion O

	1	2	3	4		108	109
F	D	A	Q	N	Q	V	Q
					L	QL
							E * *
TTTGATGCTCAAACCAAGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
31880				PstI ..		.XhoI	EcoRI

Figure 6 (Page 3 of 5)

Fusion P

	1	2	3	4		108	109
P	F	I	N	S	A	V	Q
L	Q	L	E	*	*	
CTTTTATAAACTCAGCCGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
31920			PstI ..		.XhoI		EcoRI

Fusion Q

	1	2	3	4		108	109
S	N	N	S	K	N	V	Q
L	Q	L	E	*	*	
TCAACAATTCCAAAACGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
32030		32040		PstI ..		.XhoI	EcoRI

Fusion R

	1	2	3	4		108	109
G	L	M	F	D	A	V	Q
L	Q	L	E	*	*	
GGGTTGATGTTTGACGCTGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
32030		32040		PstI ..		.XhoI	EcoRI

Fusion S

	1	2	3	4		108	109
P	N	A	P	N	T	V	Q
L	Q	L	E	*	*	
CCTAATGCACCAAACACAGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
32100			PstI ..		.XhoI		EcoRI

Fusion T

	1	2	3	4		108	109
L	E	F	D	S	N	V	Q
L	Q	L	E	*	*	
CTAGAATTTGATTCAAACGTGCAGCTGCAG...ScFv...CTCGAGTAATAAGAATTC							
32150			PstI ..		.XhoI		EcoRI

Figure 6 (Page 4 of 5)

Fusion U

		1	2	3	4		108	109
L	S	F	D	S	T	V	Q	L
						QL E * *	
CTTAGTTTTGACAGCACAGTGCAG						<u>CTGCAG</u>	...ScFv... <u>CTCGAGTAATAAGAATTC</u>	
32190					PstI ..		.XhoI	EcoRI

Fusion V

		1	2	3	4		108	109
I	D	K	L	T	L	V	Q	L
						QL E * *	
ATTGATAAGCTAACTTTGGTGCAG						<u>CTGCAG</u>	...ScFv... <u>CTCGAGTAATAAGAATTC</u>	
32240					PstI ..		.XhoI	EcoRI

Figure 6 (Page 5 of 5)

LEADHBACK AGCTAAGCTTGCATGCAAATTC
 HindIII SphI

LEADbFOR *pelB leader*
 P A M A R S Q L Q
 CCAGCGATGGCCAGATCTCAGCTGCAGAGCT
 BglIII PstI

Figure 7




Figure 8

```

          10          20          30          40          50
                                M K Y L L P
AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTATTGCCT
HindIIISphI                      SD /-----

        60          70          80          90          100          110
        T A A A G L L L L A A Q P A M A R S Q
ACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCAACCAGCGATGGCCAGATCTCAG
-----pelB leader-----| BglIIIVH

        120          130
        L Q V D G S
CTGCAGGTCGACGGATCC
PstI SalI EcoRI

```

Figure 9




Figure 10




Figure 11

TAILbBACK	<p>R S M K R A R P</p> <p>AGCTAGATCTATGAAGCGCGCAAGACCG</p> <p><i>Bgl</i>II</p>
FIBRE3FOR	<p>--fibre-----/--scFv-----</p> <p>P L N R A R Q V Q L Q</p> <p>CCTCTCAAAAAACCAAGCAGGTGCAGCTGCAGCAGCCTGG</p> <p><i>Pst</i>I</p>
FIBRE6FOR	<p>--fibre-----/--scFv-----</p> <p>P L T V H D Q V Q L Q</p> <p>CCCGCTAACCGTGCACGACCAGGTGCAGCTGCAGCAGCCTGG</p> <p><i>Pst</i>I</p>
FIBRE9FOR	<p>--fibre-----/--scFv-----</p> <p>P L T T A T Q V Q L Q</p> <p>CCTCTAACTACTGCCACTCAGGTGCAGCTGCAGCAGCCTGG</p> <p><i>Pst</i>I</p>
FIBRE12FOR	<p>--fibre-----/--scFv-----</p> <p>G V T I N N Q V Q L Q</p> <p>GGTGTGACTATTAATAATCAGGTGCAGCTGCAGGACCCTGG</p> <p><i>Pst</i>I</p>
FIBRE15FOR	<p>--fibre-----/--scFv-----</p> <p>P F D A Q N Q Q V Q L Q</p> <p>CCGTTTGATGCTCAAAACCAACAGGTGCAGCTGCAGCAGCC</p> <p><i>Pst</i>I</p>

Figure 12 (Page 1 of 2)

```

FIBRE18FOR  --fibre-----/--scFv-----
              G L M F D G Q V Q L Q
              GGGTTGATGTTTGACGCTCAGGTGCAGCTGCAGCAGCC
                                      PstI

FIBRE21FOR  --fibre-----/--scFv-----
              L S F D S T Q V Q L Q
              GCCTTAGTTTTGACAGCACACAGGTGCAGCTGCAGCAGCC
                                      PstI

FIBRE22FOR  --fibre-----/--scFv-----
              G N K N N D K L T L Q V Q L Q
              GGAAACAAAAATAATGATAAGCTAACTTTGCAGGTGCAGCTGCAGCAGCC
                                      PstI

FIBREPFOR   --fibre-----
              Y I A Q E *
              CATACATTGCCCAAGAATAACAGGTGCAGCTGCAGCAGCCTGG
                                      PstI

```

Figure 12 (Page 2 of 2)

```

          10          20          30          40          50
          M K Y L L P
AAGCTTGCATGCAAATTCTATTTCAAGGAGACAGTCATAATGAAATACCTATTGCCT
HindIII SphI                      SD /-----

          60          70          80          90          100          110
          T A A A G L L L L A A O P A M A Q V Q
ACGGCAGCCGCTGGATTGTTATTACTCGCTGCCCCAACCGCGATGGCCCCAGGTGCAG
-----pelB leader-----/VHNP

          120          130          140          150          160          170
          L Q Q P G A E L V K P G A S V K L S C
CTGCAGCAGCCTGGGGCTGAGCTTGTGAAGCCTGGGGCTTCAGTGAAGCTGTCCTGC
PstI

          180          190          200          210          220          2
          K A S G Y T F T S Y W M H W V K Q R P
AAGGCTTCTGGCTACACCTTCACCAGCTACTGGATGCCTGGGTGAAGCAGAGGCCCT
CDR1

          230          240          250          260          270          280
          G R G L E W I G R I D P N S G G T R Y
GGACGAGGCCTTGAGTGGATTGGAAGGATTGATCCTAATAGTGCTGGTACTAAGTAC
CDR2

          290          300          310          320          330          340
          N E K F L S K A T L T V D K P S S T A
AATGAGAAGTTCAAGAGCAAGGCCACACTGACTGTAGACAAACCCTCCAGCACAGCC

          350          360          370          380          390          40
          Y M Q L S S L T S E D S A V Y Y C A R
TACATGCAGCTCAGCAGCCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGA

          410          420          430          440          450
          Y D Y Y G S S Y F D Y W G Q C T T L T
TACGATTACTACGGTAGTAGCTACTTTGACTACTGGGGCCAAGGGACCACGGTCACC
CDR3 BstEII

          460          470          480          490          500          510
          V S S G G G G S G G G G S G G G G S Q
GTCTCCTCAGGTGGAGGCGGTTAGGCGGAGGTGGCTCTGGCGGTGGCGGATCCAG
/------(G4S)3 linker-----BamHI-IVλ

          520          530          540          550          560          570
          V V L T Q E S A L T T S P G E T V T L
GCTGTTGTGACTCAGGAATCTGCACTCACCACATCACCTGGTGAAACAGTCACACTC

          580          590          600          610          620
          T C R S S T G A V T T S N Y A N W V Q
ACTTGTCGCTCAAGTACTGGGGCTGTTACAACCTAGTAACCTATGCCAACTGGGTCCAA
CDR1

```

Figure 13 (Page 1 of 2)

```

630      640      650      660      670      680
E K P D H L F T G L I G G T N N R A P
GAAAAACCAGATCATTATTCACTGGTCTAATAGGTGGTACCAACAACCGAGCTCCA
                                KpnI   CDR2   SstI

      690      700      710      720      730      740
G V P A R F S G S L I G D K A A L T I
GGTGTTCCTGCCAGATTCTCAGGCTCCCTGATTGGAGACAAGGCTGCCCTCACCATC

      750      760      770      780      790      8
T G A Q T E D E A I Y F C A L W Y S N
ACAGGGGCACAGACTGAGGATGAGGCAATATATTTCTGTGCTCTATGGTACAGCAAC
                                CDR3

00      810      820      830      840      850
H W V F G G G T K L T V L G L E . .
CACTGGGTGTTCTGGTGGAGGAACCAAAGTACTGTCCTAGGTCTCGAGTAATAAGAA
                                XhoI   Eco

```

TTC
RI

Figure 13 (Page 2 of 2)




Fig. 14

(a)

1	5	10	15	20
Q V Q L Q Q S G A E L M K P G A S V K I				
MOVH-HMFG1	CAGGTT	CAGCTGCAGCAGTCTGGAGCTGAGCTGATGAGCCTCGGGCCTCAGTGAAGATA		
HUVH-HMFG1	CAGGTGCAGCTGGTGCAGTCTGGGGCAGAGGTCGAAAGCCTCGGGCCTCAGTGAAGGTC			
Q V Q L Y Q S G A E V K K P G A S V K Y				

25	30	35	40
S C K A T G Y T F S	CDR1	A Y W I E	W V K Q R
MOVH-HMFG1	TCCTGCAAGCCTACTCGCTACACATTTCAGTCCCTACTGGATAGAGTGGGTAAAGCAGAGG		
HUVH-HMFG1	TCCTGCAAGCCTTCTCGCTACACCTTCAGTCCCTACTGGATAGAGTGGGTCCGCCAGGCT		
S C K A S G Y T F S	A Y W I E	W V B O S	

45	50	52	55	CDR2
P G H G L E W I G	E I L P G S N N S R Y			
MOVH-HMFG1	CCTGGACATGGCCTTGAGTGGATTGGAGAGATTTTACCTGGAAGTAATATTCTAGATAC			
HUVH-HMFG1	CCAGGAAGGGCCTCGAGTGGGTCCGAGAGAGATTTTACCTGGAAGTAATATTCTAGATAC			
P G K G L E W Y G	E I L P G S N N S R Y			

Figure 15 (Page 1 of 4)

60	CDR2	65	70	75
MOVH-HMFG1	N E K F K G	K A T F T A D T S S N T A Y		
HUVH-HMFG1	AATGAGAAGTTCAAGGGCAAGGCCACATTCACTGCTGATACATCCTCCACACAGCCTAC	AATGAGAAGTTCAAGGGCCGAGTGACAGTCACTAGAGACACATCCACAAACACAGCCTAC		
	N E K F K G	R V T Y T B D T S I N T A Y		
80	82 A B C	85	90	95
MOVH-HMFG1	M Q L S S L T S E D S A V Y Y C S R			S Y
HUVH-HMFG1	ATGCAACTCAGCAGCCTGACATCTGAGGACTCTGCCGCTCTATTACTGTTCAAGGTCCTAC	ATGGAGCTCAGCAGCCTGAGGCTCTGAGGACACACAGCCGCTCTATTACTGTGCAAGATCCTAC		
	M E L S S L B S E D I A V Y Y C A R			S Y
	CDR3	100 A	105	110
MOVH-HMFG1	D F A W F A Y	W G Q G T P V T V S A		
HUVH-HMFG1	GACTTTGCCCTGGTTTGCTTACTGGGGCCMAGGACTCCGGTCACTGTCTCTGCA	GACTTTGCCCTGGTTTGCTTACTGGGGCCMAGGACTCTGGTCAAGTCTCTCTCA		
	D F A W F A Y	W G Q G T L V T V S S		

Figure 15 (Page 2 of 4)

(b)

1	5	10	15	20
O I V M S Q S P S S L A V S V G E K V T				
MoVx-HMFG1	GACATTGTGATGTCACAGTCTCCATCCTCCCTAGCTCTCTCAGTTGGACAGAAGGTTACT			
HuVx-HMFG1	GACATCCAGATGACCCAGAGCCCAAGCAGCCTGAGCGCCAGCGTGGGTGACACAGAGTGACC			
D I Q M I Q S P S S L S A S V G Q R V T				
	25	27	A B C D E F	30
M S C	K S S Q S L L Y S S N Q K I Y L A			CDRL1
MoVx-HMFG1	ATGAGCTGCAAGTCCAGTCAGAGCCTTTTATATAGTAGCAATCAAAAGATCTACTTGGCC			
HuVx-HMFG1	ATCACCTGTAAAGTCCAGTCAGAGCCTTTTATATAGTAGCAATCAAAAGATCTACTTGGCC			
I T C	K S S Q S L L Y S S N Q K I Y L A			
35	40	45	50	CDR2
W Y Q Q K P G Q S P K L L I Y			W A S T R	
MoVx-HMFG1	TGGTACCAGCAGAAACCAGGCCAGTCTCCTAACTGCTGATTTACTGGGCATCCACTAGG			
HuVx-HMFG1	TGGTACCAGCAGAAAGCCAGGTAAAGCTCCAAAGCTGCTGATCTACTGGGCATCCACTAGG			
W Y Q Q K P G K L P K L L I Y			W A S T R	

Figure 15 (Page 3 of 4)

Figure 15 (Page 4 of 4)